

Dr. Bbosa Science

This document is sponsored by

The Science Foundation college Kiwanga-Namanve,
Uganda East Africa

Senior one to Senior six,

+256 778 633 682, +256 753 802709

Based on sciences, Best for Sciences

Obtain free notes, tests, making guides from: digitalteachers.co.ug

Respiration

This is the oxidation of organic substance to liberate energy in the body. **Aerobic respiration** requires oxygen whereas **anaerobic respiration** does not require oxygen.

Organic molecules (usually carbohydrate or fat) are broken down bond by bond, by a series of enzyme - controlled reactions. Each bond broken releases a small amount of energy converted to **adenosine triphosphate (ATP)**. ATP is the immediate source of energy for cellular reactions.

Uses of ATP

1. Provide energy to build up macromolecules such as proteins from amino acids, polysaccharides from monosaccharides and DNA synthesis.
2. Provide energy for movement of materials such as active transport.
3. Provides energy for secretion of materials.
4. Provide energy for muscle contraction, spindle formation in cell division and ciliary action
5. Provide energy for activation of molecules before they are used in the body.

Cell respiration.

Cell respiration involves oxidation of a substrate to yield chemical energy (ATP). Organic compounds which are used as substrates in respiration are carbohydrates, fats and proteins.

Carbohydrates:

These are usually the first choice of most cells. In fact, brain cells of mammals cannot use anything but glucose. Polysaccharides are hydrolyzed to monosaccharides before they enter the respiratory pathway i.e. starch in plant and glycogen in animals are first converted to glucose.

Fats/lipids.

They form the “first” reserve and are mainly used when carbohydrate reserves have been exhausted. However, in skeleton muscle cells, if glucose and fatty acids are available, these cells respire the acids in preference to glucose.

Lipids are better energy source than carbohydrates because they have a higher proportion of hydrogen and an almost insignificant proportion of oxygen compared with carbohydrates. Thus a given mass of lipids yields more energy on oxidation than an equal mass of carbohydrate.

Proteins.


Since proteins have other essential functions they are only used when all carbohydrate and fat reserves have been used up, as during prolonged starvation.

The break of glucose in respiration.

When glucose is the substrate of respiration, its oxidation can be divided into three phases, glycolysis, oxidative de-carboxylation (Krebs or citric acid cycle or TCA, tri-carboxylic acid cycle); oxidative phosphorylation. Glycolysis is common to anaerobic and aerobic respiration, but the other two phases only occur only in presence of oxygen.

Stages in glycolysis in the cytoplasm of the cell

1. Glucose (6C sugar)




The glucose molecule is phosphorylated to make it more reactive. The phosphate molecule comes from conversion of ATP to ADP

2. Glucose phosphate(6C)



Glucose is reorganized into its isomer fructose phosphate

3. Fructose




Further phosphorylation takes place by the addition of another phosphate group from ATP make the sugar more active

4. Fructose bisphosphate



Sugar splits into two 3 carbon sugars

5. Glyceraldehyde-3-phosphate
(2molecules)




Further phosphorylation takes place by the addition of another phosphate group from inorganic sources but not ATP. 2pairs of hydrogen removed

6. Glycerate-1,3-bisphosphate
(2molecules)



A phosphate molecule is lost from each glycerate-1,3-bisphosphate molecule yielding two ATP molecules from two ADP molecules.

7. Glycerate-3-phosphate
(2molecules)



A phosphate is removed from the each glycerate-3-phosphate to form a pair of ATP

8. Pyruvate (3C)
(2molecules)

Glycolysis represents a series of reactions in which a glucose molecule is broken down into two molecules of pyruvate. Glycolysis occurs in the **cytoplasm of cells**, not in the mitochondria, and does not require the presence of oxygen. The process may be subdivided into two steps, first the conversion of glucose into fructose 1,6 - di-phosphate, and secondly the splitting of fructose - 1,6- di-phosphate into 3C sugars which are later converted into pyruvate. Two ATP molecules are used up for phosphorylation reaction in the first step. Whilst four ATP molecules are produced in the second step. Four hydrogen atoms are also release. These are oxidized to water by molecular oxygen with accompanying phosphorylation of ADP to ATP molecules during oxidative. Each pair of hydrogen atom is oxidized to form 2 ATP

The ultimate fate of pyruvate depends on the availability of oxygen in the cell. If it is present, pyruvate will enter a mitochondrion and be completely oxidized into carbon dioxide and water **aerobic respiration**. If oxygen is unavailable pyruvate will be converted into ethanol or lactate (**anaerobic respiration**).

Summary

source	Number of ATP
From stages 6 and 7	4ATP
ATP from NADH ₂	4ATP
Total	8ATP

Aerobic respiration.

There are two phases involved in aerobic respiration, first, if sufficient oxygen is available, each pyruvate molecule enters a mitochondrion where it is converted into a 2-carbon compound **acetyl co-enzyme A** (acetyl CoA for short). In this reaction carbon dioxide is given off, and the pyruvate loses a pair of hydrogen atoms which again results in the synthesis of ATP.


Acetyl CoA is a very important intermediate in respiration. It links glycolysis, oxidation of fats and proteins with the **Krebs cycle**.

The Krebs cycle, tricarboxylic cycles or citric cycle

1. Before pyruvate enters the Krebs cycle it combines with a compound called **coenzyme A** to form Acetyl coenzyme A. in the process, a molecule of carbon dioxide and a pair of hydrogen atoms are removed.
2. The 2-carbon acetyl coenzyme A enters the Krebs cycle by combining with a 4-carbon **oxaloacetate (oxaloacetic acid)** to give a 6-carbon **citrate (citric acid)**. This reaction requires energy which is provided at the expense of the energy-rich bond of acetyl group CoA.
3. The citrate is degraded to a 5-carbon **α -ketoglutarate (α -ketoglutaric acid)** and then the 4-carbon oxaloacetate by progressive loss of two carbon dioxide thus completing the cycle.
4. For each turn of the cycle, a total of four pairs of hydrogen atoms are also formed.

- Of these, 3 pairs of hydrogen atoms are combined with hydrogen carrier, nicotinamide adenine dinucleotide (NAD) and yield three ATP for each pair of hydrogen atoms. The remaining pair of hydrogen atoms combine with a different hydrogen carrier, **Flavine adenine dinucleotide (FAD)**. And yield 2 ATP.
- It must be remembered that all these products are formed from a single pyruvate molecule of which two are produced from each glucose molecule.

Krebs Cycle in the matrix of the mitochondria




Summary

Source of ATP	Number of ATP	Total number
3 pairs of H with NAD	9ATP	9ATP x 2
1 pair of H with FAD	2ATP	2ATP x 2
Energy of the cycle	1ATP	1ATP x 2
Glycolysis process	8ATP	8ATP x 1
Pyruvate to Acetyl Co A	3ATP	3ATP x 2
Total		38 ATP


Electron transport in the stalked membrane of the mitochondria

It occurs in the membranes of the mitochondria. It is the means in by which the energy, in the form of hydrogen atom, from Krebs cycle, is converted to ATP. The hydrogen atoms attached to a hydrogen carriers NAD and FAD are transferred to a chain of other carriers at progressively lower energy levels. As the hydrogens pass from one carrier to the next, the energy released is used to produce ATP. The series of carrier is called **respiration chain**. The carriers in the chain include **NAD, Flavo protein, coenzyme Q** and iron-containing protein called **cytochromes**. Initially hydrogen atoms are passed along the chain, but latter split into their proton and electron and only the electron pass from carrier to carrier. For this reason, the pathway can be called electron or hydrogen transport system. At the end of the chain the protons and electrons recombine, and the hydrogen atoms create a link with oxygen to form water. This formation of ATP through the oxidation of the hydrogen atoms is called **oxidative phosphorylation**. The role of oxygen is to act as the final electron acceptor.

Summary of the electron transport system



Mitochondrion



Function

Production of cellular energy, ATP

Adaptations

- Has extensive inner membrane to create a high surface area for enzymes that produce.
- have a small amount of mitochondrial DNA, allowing them to create mitochondrial proteins quicker than from nucleic genes.
- Have necessary enzymes for production of energy

Anaerobic respiration

A variety of microorganism (anaerobes) employ anaerobic respiration as their major ATP yielding process. Organism that survive only in absence of oxygen are termed **obligate anaerobes** e.g. *C. Botulium* and *C. tetani*). Obligate anaerobe find oxygen poisonous.

Other organisms such as yeast and alimentary canal parasites (such as tape worms), can exist whether oxygen is available or not. These are called **facultative anaerobe**. Also, some cells that are temporarily deprived of with **no oxygen available to accept the hydrogen** oxygen (such as muscle cells) are able to respire anaerobically.

Pyruvate serves as an electron/hydrogen acceptor in absence of oxygen to accept the hydrogen atoms released during glycolysis; and depending on the metabolic pathways within the organisms' cells, the end-product of anaerobic respiration will either be ethanol and carbon dioxide (e.g. fermentation as in yeast) or lactate, for example **lactate fermentation** in muscle cells:

In Alcoholic fermentation the glucose is converted to ethanol and carbon dioxide



Alcoholic fermentation is the basis of brewing in which ethanol is an important product and baking industry in which carbon dioxide expands the dough.

Lactic fermentation occurs occasionally in animal cells during strenuous exercise and oxygen is insufficient. It allows animal to survive periods of insufficient oxygen. When oxygen is latter availed, lactic acid is oxidized to carbon dioxide and water or can be turned into carbohydrates. The amount of oxygen required to oxidize lactic acid accumulated in muscles is called the **oxygen debt**.

Differences in aerobic and anaerobic respiration

	Aerobic Respiration	Anaerobic Respiration
Definition	Aerobic respiration uses oxygen.	Anaerobic respiration is respiration without oxygen; the process uses a respiratory electron transport chain but does not use oxygen as the electron acceptors.
Cells that use it	Aerobic respiration occurs in most cells.	Anaerobic respiration occurs mostly in prokaryotes
Amount of energy released	High (36-38 ATP molecules)	Lower (Between 36-2 ATP molecules)
Stages	Glycolysis, Krebs cycle, Electron Transport Chain	Glycolysis, Krebs cycle, Electron Transport Chain
Products	Carbon dioxide, water, ATP	Carbon dioxide, reduced species, ATP
Site of reactions	Cytoplasm and mitochondria	Cytoplasm and mitochondria
Reactants	glucose, oxygen	glucose, electron acceptor (not oxygen)
combustion	complete	incomplete
Production of Ethanol or Lactic Acid	Does not produce ethanol or lactic acid	Produce ethanol or lactic acid

Alternative respiratory substances

1. **Fats** are oxidized after hydrolysis to glycerol and fatty acids. Glycerol is phosphorylated, and converted into glyceraldehyde-3-phosphate which is incorporated into the glycolysis pathway. Fats produce more hydrogens than glucose/carbohydrates and hence produce more energy.
2. **Proteins** are used only in cases of starvation. They are hydrolyzed to constituent amino acids which are deaminated to form carbohydrates before converted to pyruvate.

Respiratory Quotients

The respiratory quotient (RQ) is the measure of the ratio of carbon dioxide evolved by an organism to the oxygen consumed, over a certain period.

$$\text{RQ} = \frac{\text{CO}_2 \text{ evolved}}{\text{O}_2 \text{ consumed}}$$

For hexose sugar like glucose, the equation for its complete oxidation is



$$\text{RQ} = \frac{6\text{CO}_2}{6\text{O}_2} = 1.0$$

The **respiratory quotient** (RQ) reflects the relative contributions of fat, carbohydrate, and protein to the oxidation fuel mixture.

In fats and proteins, the ratio of carbon dioxide evolved to oxygen consumed is far smaller than in carbohydrates. A fat and proteins, therefore requires more oxygen for its complete oxidation than carbohydrates. The respiratory quotient of fats is 0.7 whereas that of proteins is 0.9.

Organisms rarely respire one substance at a time, thus the respiratory quotients of an animal is between 0.8 and 0.9.

Low respiratory quotient may also indicate that carbon dioxide produced is used elsewhere for example;

- (i) at compensation point during photosynthesis carbon dioxide produced used is in photosynthesis
- (ii) during egg formation carbon dioxide produced is used for shell formation.

High respiratory quotients occur when carbohydrates are converted into fat e.g. when an animal is preparing to hibernate and during anaerobic respiration because carbon dioxide is given out but no oxygen used.

Basal metabolic rate

Is the minimal rate of energy expenditure per unit time by endothermic animals at rest.

Factors affecting the BMR

1. **Body size:** Metabolic rate increases as weight, height, and surface area increase.
2. **Body composition:** Fat tissue has a lower metabolic activity than muscle tissue. As lean muscle mass increases, metabolic rate increases.

3. **Gender:** The basal metabolic rate (BMR) averages 5 to 10 percent lower in women than in men. This is largely because women generally possess more body fat and less muscle mass than men of similar size.
4. **Age:** A decrease in lean muscle mass during adulthood results in a slow, steady decline of roughly 0.3 percent per year in BMR after the age of about 30. This can be largely avoided by strength training throughout adulthood.
5. **Climate and body temperature:** The BMR of people in tropical climates is generally 5 to 20 percent higher than their counterparts living in more temperate areas because it takes energy to keep the body cool. Exercise performed in hot weather also imposes an additional metabolic load. Body fat content and effectiveness of clothing determine the magnitude of increase in energy metabolism in cold environments; it takes energy to keep the body warm if you work or exercise in very cold weather.
6. **Hormonal levels:** Thyroxine (T₄), the key hormone released by the thyroid glands has a significant effect upon metabolic rate. Hypothyroidism is relatively common, especially in women near or after menopause. Everyone with a weight problem should have their thyroid function checked by their doctor and treated appropriately if it turns out to be low.
7. **Health:** Fever, illness, or injury may increase resting metabolic rate two-fold


Exercise


Section A

1.	Lactic acid accumulation in muscles of an athletic during action is due to A. Oxygen debt B. Anaerobic respiration C. Panting D. High rate of respiration
2.	The end product of glycolysis A. Glucose diphosphate B. Lactic acid C. Citric acid D. Pyruvic acid
3.	The following process does not require respiratory energy? A. Synthesis of cellulose B. Meiosis C. Loss of water from the stomata D. Mineral absorption
4.	Which of the following are formed during anaerobic respiration in yeast cell? A. Lactic acid and ATP B. Lactic acid and ADP C. Ethanol and ATP D. Ethanol and ADP
5.	How many kilojoules of energy are released if 1 gram of sugar burned in oxygen raises the temperature of 500g of water by 7.5°C , (4.18J raise the temperature of 1 gram of water by 1°C) A. 15.7kJ B. 156kJ C. 1.56kJ D. 1560kJ
6.	What happens to most of the reduced NADH ₂ molecules metabolism? A. Direct use in the synthesis of starch to glucose B. Oxidation in mitochondria resulting in ATP formation C. Oxidation in Calvin cycle D. Combination with sulphuric acid as part of Krebs cycle
7.	Which of the following will speed of phosphorylation of the Hexose sugar? A. Decrease of ADP concentration B. A decrease in the concentration of ATP C. An increase in the concentration of the phosphate D. An increase in the concentration of phosphorylated hexose

8.	<p>An athletic had just finished a race; the phrase “oxygen debt” refers to</p> <ul style="list-style-type: none"> A. The amount of oxygen originally present in the muscle of an athlete before the race B. The total amount of oxygen an athlete requires to restore the breathing rate to normal C. The amount of oxygen taken in after the race and used for complete combustion of lactic acid D. The amount of oxygen after the race to convert excess lactic acid to glycogen in the liver.
9.	<p>Which of the following biological processes does not utilize respiratory energy?</p> <ul style="list-style-type: none"> A. Loss of water from stoma B. Mineral salt absorption C. Synthesis of cellulose D. Meiosis
10.	<p>Which one of the following compounds act as hydrogen acceptor during anaerobic respiration in animals?</p> <ul style="list-style-type: none"> A. NAD B. NADP C. Lactic acid D. Pyruvic acid
11.	<p>Which one of the following is not a method of measuring the rate of respiration in an organism?</p> <ul style="list-style-type: none"> A. Estimating the amount of food taken in by the organism per day B. Measuring the heat produced by the organism in a given time C. Measuring the amount of CO₂ produced by an organism in a given time D. Estimating the amount of oxygen consumed by the body in a given time
12.	<p>The substance that supplies phosphate at the beginning of glycolysis is</p> <ul style="list-style-type: none"> A. Adenosine diphosphate (ADP) B. Adenosine triphosphate (ATP) C. Adenosine monophosphate (AMP) D. Nicotinamide adenine dinucleotide phosphate (NADP)
13.	<p>The following are physiological conditions in living cells</p> <ol style="list-style-type: none"> 1. High concentration of ADP and Pi 2. High concentration ATP 3. High concentration of hydrogenase 4. High concentration of ATPase <p>Which of them will increase the rate of sugar down</p> <ul style="list-style-type: none"> A. 1 and 2 B. 2 and 3 C. 1 only D. 4 only

14.	<p>A rat requires more energy per unit body weight than required by human because the rat</p> <ul style="list-style-type: none"> A. Has a large surface area B. Is more active C. Has higher body temperature D. Has a higher metabolic rate
15.	<p>In which of the following does anaerobic respiratory not occurs?</p> <ul style="list-style-type: none"> A. Skeletal muscle B. Yeast cell C. Bacterial D. Smooth muscle
16.	<p>The life process which releases most energy is</p> <ul style="list-style-type: none"> A. The light reaction of photosynthesis B. Fermentation of glucose C. Aerobic cellular respiration of glucose D. The oxidation of lactic in mammalian of muscle cells
17.	<p>The respiration of the fat tri olein can be summarized as the follows by the equation</p> $C_{57}H_{104}O_6 + 80O_2 \rightarrow 57CO_2 + 52H_2O$ <p>Which of the following resents the respiratory quotient (RQ) for this fat?</p> <ul style="list-style-type: none"> A. 1.5 B. 1.4 C. 0.71 D. 0.65
18.	<p>Which biological process takes place in the mitochondrion?</p> <ul style="list-style-type: none"> A. Glycolysis B. Formation of lactic acid C. Tricarboxylic cycle D. Alcoholic fermentation
19.	<p>Which of the following conversion takes place in human under conditions of starvation?</p> <ul style="list-style-type: none"> A. Fatty acids to carbohydrate B. Proteins to carbohydrates C. Glucose to lipids D. Lipids to lipoproteins
20.	<p>During which one of the following is the respiratory quotient most likely to be high</p> <ul style="list-style-type: none"> A. In plants during bright light B. In animals during laying down of fats C. During egg laying in birds D. During lactic acid formation

21.	<p>The figure below shows some reactions in the respiratory pathway</p> <div style="text-align: center;">  </div> <p>The enzyme which catalyzes the reaction marked X is</p> <p>A. Pyruvate dehydrogenase B. Acetyl CoA dehydrogenase C. Pyruvate decarboxylase D. Acetyl CoA decarboxylase</p>
22.	<p>During strenuous activity, the pyruvic acid in the muscle may accept hydrogen from reduced NAD to become</p> <p>A. Acetyl CoA B. Lactic acid C. Ethanol D. Citric acid</p>
23.	<p>The equation for complete oxidation of a substance is</p> $2C_{18}H_{34}O_2 + 51O_2 \longrightarrow 36CO_2 + 34H_2O$ <p>The respiratory quotient for oxidation is</p> <p>A. 0.70 B. 1.4 C. 0.9 D. 1.0</p>
24.	<p>Which of the following is liberated during both aerobic and anaerobic respiration?</p> <p>A. Carbon dioxide and energy B. Ethanol and water C. Water and carbon dioxide D. Carbon dioxide and ethanol</p>
25.	<p>In endergonic reaction, the products of the reaction contain</p> <p>A. More energy than the reactants and energy is released B. Less energy than the reactants and energy is absorbed C. More energy than the reactants and energy is supplied D. Less energy than the reactants and energy is released</p>
26.	<p>Which of the following is the ultimate hydrogen acceptor during anaerobic respiration?</p> <p>A. Lactic acid B. NAD C. Pyruvic acid D. Acetaldehyde</p>

27.	<p>Which one of the following is unlikely to be found in the body of obligate anaerobes?</p> <p>A. Glycolytic enzymes B. ATP C. Mitochondria D. Acetaldehyde</p>
28.	<p>The figure shows that</p>  <p>A. NAD is oxidized to NADH₂ B. NADH₂ is reduces FAD to FADH₂ C. FADH₂ is reduced is reduced to FAD D. NADH₂ + H₂ → NAD</p>
29.	<p>Anaerobes thrive better than aerobic organism in water experiencing thermal pollution because</p> <p>A. High temperatures kill aerobic organisms B. Anaerobes possess enzymes that work best at high temperatures C. High temperatures reduce solubility of oxygen D. High temperatures encourage multiplication of aerobes predators.</p>
30.	<p>A major difference between respiration and burning is that</p> <p>A. no heat is produced during respiration B. burning is a faster process C. burning is a chemical process D. chemical energy is stored in respiration</p>
31.	<p>Compared to carbohydrates, fats have high energy value because fats</p> <p>A. have long chains of fatty acids B. have a higher proportional of hydrogen C. are more compact in structure D. have a high proportion of oxygen</p>


32.	<p>The compound which acts as oxidizing agent during anaerobic respiration in plants</p> <p>A. NAD and pyruvate B. Ethanal and NAD C. NAD and FAD D. NADP and pyruvic acid</p>
33.	<p>Which of the following increases the rate photophosphorylation of hexose sugar during the normal respiration process?</p> <p>A. An increase in ADP concentration B. An increase in ATP concentration C. An increase in concentration of hexose sugar D. A decrease in concentration of phosphorylated sugar</p>
34.	<p>Which one of the following activities in living organisms can result in a respiratory quotient of less than 1.0?</p> <p>A. When carbohydrates are respired B. During extensive laying, down of fat in livestock C. At compensation point, during photosynthesis D. When the rate of exhalation equals that of inhalation.</p>
35.	<p>A rat requires more energy per unit body weight than that required by a human because a rat</p> <p>A. Has a larger surface area B. Is more active C. Has higher body temperature D. Has a higher metabolic rate</p>
36.	<p>Lipids are better energy sources than carbohydrates is that</p> <p>A. Are insoluble B. Do not form hydrogen bonds with water C. Are more compact D. Have a higher proportion of hydrogen</p>
37	<p>Which one of the following changes occur in mammalian body at the onset of an exercise?</p> <p>A. Increase in the pH of blood B. Decrease in the rate of contraction of the diaphragm muscles C. Increase in the rate of tissue respiration D. Decrease in the amounts of water vapor in the breath</p>
38	<p>Fats yield more energy per unit mass than carbohydrates because fats possess</p> <p>A. More carbon atoms B. More hydrogen atoms C. Few carbon atoms D. Fewer oxygen atoms</p>
39	<p>Which one of the following best describes basal metabolic rate?</p> <p>A. Average amount of energy produced by the body B. Average amount of energy produced when at rest C. Amount of energy produced by an average body D. Amount of energy produced when all voluntary movement have</p>

	ceased.
40	Lactic acid accumulation in the muscle of an athlete during action is due to A. Oxygen debt B. Anaerobic respiration C. Panting D. High rate of respiration
41	In which one of the following parts of the cell does most production of ATP occurs? A. Matrix of mitochondria B. Cristae of the mitochondria C. Cytoplasm of the cell D. Outer membrane of the mitochondria
42	Which one of the following is not a method of measuring the rate of respiration in an organism? A. Estimating the amount of food taken in by the organism per day B. Measuring the heat produced by the organism in a given time C. Measuring the amount of carbon dioxide produced by an organism in a given time D. Estimating the amount of oxygen consumed in a given time
43	In an individual whose energy intake is 50000kJ per day, loses 1dm ³ of sweat in a day during manual work. Given that the latent heat of vaporization of sweat is 2.45kJcm ⁻³ , the percentage of energy lost by the individual during the work is A. 4.22 B. 5.0 C. 4.9 D. 7.65
44.	The latent heat of evaporation of sweat is 3.15kJcm ⁻³ . What is the percentage of energy lost by sweating from a manual worker who loses 2dm ³ per day of sweat and has a daily energy intake of 40,000kJ? A. 6.30 B. 7.88 C. 8.25 D. 15.75
45	In which of the following does anaerobic respiration not occur A. Skeletal muscles B. Yeast cell C. Bacteria D. Smooth muscle
46	Which one of the following food materials has the highest amount of potential energy per unit weight? A. Proteins B. Vitamins C. Monosaccharide D. Fats

47 Which one of these activities would result into a low respiratory quotient?

- A. Respiration in muscles during heavy exercise
- B. Formation of calcareous shells
- C. Fattening livestock
- D. Preparation for hibernation in a mammal.

48. The equation for respiration of a substrate is



What is the respiratory quotient of substrate?

- A. 0.70 B. 0.80 C. 0.90 D. 1.0

49. During respiration in the absence of oxygen, pyruvic acid is converted

- A. lactic acid and water
- B. ethanol and carbon dioxide in plants
- C. lactic acid and carbon dioxide in plants
- D. ethanol and water in plants


Structured questions

50. Fill in the missing using appropriate words (s) so as to complete the account of cellular respiration

‘the initials stage in cellular respiration is glycolysis which occurs with occur in the of the cell. A monosaccharide such as And is first Then converted into a 3-carbon intermediate product called

In the absence of oxygen this product may enter another metabolic pathway giving rise to in animals and in plants. The end product of glycolysis is Which is decarboxylated and enter the Cycle. This cycle links other pathways which enables organism acid such as and To be regarded and small organic molecules in the cycle provides uses into the reduction of during cycle a total of molecules are formed from complete oxidation per molecule of a 6-carbon sugar

51.



- (a) Where exactly do reactions P, Q and R occur?
 P
 Q
 R
- (b) What is substance Z?

- (c) What is the 6C intermediate compound from the reaction between acetyl CoA and the 4C compound
- (d) Name the enzyme controlling reaction Y

- (e) Name the final hydrogen acceptor in the cytochromes system indicated in the diagram

- (f) (i) Define the term respiratory Quotient (RQ)

- (ii) Glucose has a formula $C_6H_{12}O_6$ and tripalmitin has the formula $C_{51}H_{99}O_6$. The two substances when completely oxidized have the following overall equation.



Using the above equations calculate RQ values for aerobic respiration of tripalmitin

- (iii) State the RQ value of glucose in anaerobic respiration. Explain your answer.

52. (a) Differentiate between respiratory quotient (RQ) and basal metabolic rate (BMR)

- (b) The table below shows the respiratory quotient in germinating seeds under different treatment

	Treatment	RQ
(i)	4hrs soaking in water	6.0
(ii)	4hr soaking then 4 hour exposure to air	1.8
(iii)	4hour soaking the 24hrs exposure to air	1.0

Explain the difference respiratory quotients of the germinating seeds under the different treatment. (06 marks)

- (a) Explain why the BMR varies with age of individual (02marks)


54. (a) Name the organelle most important in photosynthesis and respiration
 (i) Photosynthesis
 (ii) Respiration
 (b) Give four differences and five similarities between the two organelles
 (i) Differences
 (ii) Similarities
55. (a) Give an account of glycolysis
 (b) Explain what happened to the pyruvate under aerobic conditions.
56. (a) Distinguish between aerobic respiration and anaerobic respiration
 (b) Outline the major steps involved in the process of glycolysis in a cell.
 (c) How is energy produced in glycolysis?
57. (a) Differentiate between aerobic and anaerobic respiration. (05marks)
 (b) Describe what happens to the end product of glycolysis in absence of oxygen (10marks)
 (c) Why is it important to produce ATP during cellular respiration? (05marks)
58. An investigation was carried out into the effects of athlete training on respiration in human muscle. A group of athlete and non-athlete exercised at different levels. The level of exercise was expressed as rates of energy expenditure $\text{Jmin}^{-1}\text{kg}^{-1}$ of body mass. At each level, rates of oxygen consumption and lactic acid production were measured, the results are shown in the table below. Figures in the table are the means of the measurements made for each group.

Rate of energy Expenditure in $\text{Jmin}^{-1}\text{kg}^{-1}$	Rate of oxygen consumption in $\text{cm}^3\text{min}^{-1}\text{kg}^{-1}$		Rate of lactic acid production in $\text{mgmin}^{-1}\text{kg}^{-1}$	
	Athletes	Non-athletes	Athletes	Non-athletes
600	30	29	0	0
800	40	39	0	0
1000	50	44	0	185
1200	57	45	85	350
1400	58	45	305	590

- (a) Plot graphs to show the relationship between the rate of oxygen consumption and lactic acid production against the level of activity in the two groups.
 (b) Using your graphs in (a) describe the effect of increasing the level of exercise on the rates of oxygen consumption and lactic acid accumulation in
 (i) Athletes
 (ii) Non-athletes
 (c) How do the two groups of people compare in their abilities to consume oxygen and accumulate lactic acid during the exercise?
 (d) Using the data provided, suggest any advantage an athlete has over a non-athletes.

59. An experiment was carried out to investigate the rate of respiration of yeast cells mixed with three different carbohydrates (glucose, sucrose and lactose), using methylene blue as an indicator (methylene blue is blue in alkaline condition and colorless in acidic condition).

1cm³ of 0.1M methylene blue was added to a mixture of 5cm³ of a suspension of yeast in 10cm³ of 0.5% glucose solution in boiling tube. The boiling tube was placed in a water bath at 30⁰C for 20 minutes. The rate of respiration was measured as a percentage of the intensity of the blue color at the beginning of the experiment, at interval of 2minutes. The experiment was repeated using 5% sucrose and lactose. The results are shown in the figure below. Study the figure and answer the questions that follow.



- Calculate the average rate of respiration of yeast in glucose solution during the first four minutes in terms percentage intensity of blue color. (3minutes)
- Describe the change in the intensity of blue color with time, for each carbohydrate. (05marks)
- Explain the relationship described in (b) for each carbohydrate,
 - Lactose (03marks)
 - Sucrose
 - glucose
- suggest what would happen to the color for glucose and sucrose if the experiment continued for 10 more minutes. Give an explanation in each case (10marks)
- Explain why the boiling tubes were
 - Kept covered during the experiment. (03marks)

(ii) Placed in a water bath at 30⁰C. (3marks)

		Paper 2
11.	1997/2/6	(a)
12.	1995/2/3	(a)
13.	2015/2/4	
14.	2000.2.1	(e)
15	2013/2/1	

Answers to objective question

1.	B	11.	A	21.	A	31.	B	41.	B
2.	D	12.	B	22.	B	32.	A	42.	A
3.	C	13.	C	23.	A	33.	D	43.	C
4.	C	14.	A	24.	C	34.	C	44.	D
5.	A	15.	D	25.	C	35.	A	45.	D
6.	B	16.	C	26.	C	36.	D	46.	
7.	B	17.	c	27.	C	37.	C	47.	
8.	C	18.	C	28.	B	38.	B	48.	
9.	A	19.	B	29.	C	39.	B	49.	
10.	D	20.	D	30.	D	40.	B		

50. Cytoplasm, glucose, fructose, phosphorylated, lactic acid, ethanol, pyruvic acid, Kreb's cycle, fatty acid, amino acid, hydrogen atoms, oxygen, 38.

51. (a) P- glycolysis

Q – Kreb's cycle

R- electron transport system

(b) Z – carbon dioxide

(c) citrate

(d) decarboxylase

(e) oxygen

(f)(i) respiratory quotient = $\frac{\text{carbon dioxide given out}}{\text{oxygen used}}$

$$(ii) RQ = \frac{51}{\left(\frac{145}{2}\right)} = 0.7$$

(iii) Infinity because carbon dioxide is given out without use of oxygen

(c) Solution

(a) Respiratory quotient is the ratio of the volume of carbon dioxide given out to the volume of oxygen consumed when one mole of respiratory substrate is completely oxidized in the body.

52. (a) Basal metabolic rate is the rate at which the minimum amount of energy needed to maintain vital processes of the body cells such as temperature regulation and breathing, is released when the body is completely at rest.

(b) When the seeds are not exposed to air, there is inadequate oxygen supply and absorption.

The seeds respire anaerobically to produce carbon dioxide.

The volume of carbon dioxide released by the seeds in respiration far exceeds that of oxygen consumed and the RQ is greater than 1.0.

When exposed to air for 4hrs, the volume of oxygen absorbed by the seeds increases. Then, there is a combination of aerobic and anaerobic respiration occurring. However, the volume of carbon dioxide produced exceeds that of oxygen consumed and the RQ is still above 1.0 but less than in (i) above.

When exposed to air for 24hrs, the volume of oxygen absorbed is adequate for the plant to rely solely on aerobic respiration for its energy needs.

Since carbohydrates are the main energy substrate in the seed, complete respiration consumes the same volume of oxygen as the volume of carbon dioxide produced and so the RQ is 1.0.

(c) At rest, the energy requirements of the body vary with age to keep pace with the energy requirement at a particular age, BMR also varies with age.

53. Photosynthesis – chloroplast

Respiration - mitochondria

Similarities of Mitochondria and Chloroplasts

- Produce energy
- have its own DNA
- enclosed by two membranes
- oxygen (O₂) and carbon dioxide (CO₂) are involved in its processes
- both have fluids inside of them

Differences Between Mitochondria and Chloroplasts

	Chloroplast	Mitochondria
1.	They are usually found in plants	They are found in plants and animal cell
2.	They convert light energy into chemical energy (sugar)	Breakdown chemical energy (sugar) into another form of chemical energy (ATP)
3.	Contain chlorophyll	Lack chlorophyll
4.	Inner membrane is thylakoid	Inner membrane is cristae

58 (a)

(b) (i) Athlete

- An increase in the level of activity leads to steady/gradual/uniform rise in the rate of oxygen consumption up to the level of 1000 after which the rate of increase is slower and tend to level off.
- Lactic acid accumulation increases exponentially after energy consumption goes beyond $1000\text{JMin}^{-1}\text{kg}^{-1}$.

(ii) non-athletic

- An increase in the level of activity leads to steady/gradual/uniform rise in the rate of oxygen consumption up to the level of 1000 after which the rate of increase is slower and tends to level off.
- Lactic acid accumulation increases exponentially after energy consumption goes beyond $800\text{JMin}^{-1}\text{kg}^{-1}$.

(c)

- Athletes consume oxygen at a higher rate than non-athletes
- Non-athlete produce more lactic acid than athletes
- Non athlete starts to produce lactic acid at lower rate of energy consumption.

(d)

Athletes consume more oxygen leading to complete respiration producing more energy than non-athlete. They thus produce less lactic acid.

60. Solution:

$$\begin{aligned} \text{(a) Rate of respiration} &= \frac{\text{change in percentage intensity of blue colour}}{\text{time taken}} \\ &= \frac{100-52}{4} \\ &= \frac{48}{4} \end{aligned}$$

Rate of respiration = 12% per minute

Hence the average rate of respiration of yeast in glucose solution during the first four minutes is 12% per minute

(b)

- Initially, the percentage intensity of blue colour is 100% for all carbohydrates.

Lactose

- The percentage intensity of blue colour remains the same throughout the time of the experiment.

Sucrose

- The percentage intensity of blue decreases gradually and linearly with time

Glucose

- The percentage intensity of blue colour decreases rapidly up to the 8th minute. It then decreases gradually up to the 18th minute after which it remains more or less constant up to the end of the experiment.

(c) (i) lactose

Lactose is not a substrate of yeast metabolism and therefore is not utilized throughout the experiment. This may be as a result of lack of enzyme, lactase, in the yeast cells to hydrolyze lactose into glucose and galactose.

As a result, the pH of the contents of the reaction medium remains the same throughout the experiment.

(ii) Sucrose

Sucrose is readily hydrolyzed into glucose and fructose by the enzyme, invertase, present in yeast. Glucose then undergoes anaerobic respiration to produce alcohol and carbon dioxide is an acidic gas and therefore lowers the pH of the contents of the boiling tube and so reduces the percentage of the intensity of the blue colour.

Sucrose acts as a ready stock of glucose which is increasingly broken down to provide glucose for respiration of the yeast cells. As a result, increasingly more carbon dioxide is produced, thereby progressively increasing the acidity of the medium and therefore reducing the percentage intensity of the blue colour continually.

(iii) Glucose

Initially, the concentration of glucose is high. This is utilized directly by the yeast cells, respiring anaerobically to produce carbon dioxide and alcohol. Since the glucose is readily available, the rate of production of carbon dioxide is higher for glucose than sucrose and therefore reduces the percentage intensity of the blue colour more rapidly with time.

Decreasing rate of reduction in percentage intensity of the blue colour occurs because;

- The amount of glucose remaining in the boiling tube has greatly.
- The yeast cells have started to be inhibited/ killed by the accumulation of ethanol in the reaction medium.

As a result, the rate of respiration decreases progressively at a decreasing rate until equilibrium is attained and it remains constant.

(d) The colour for glucose would remain the same throughout the extra 10 minutes.

- This is because the yeast cells are inhibited by the accumulation of ethanol so that further respiration of the available glucose, if any, does not continue. Therefore, the pH of the medium remains the same for the entire extra 10

The colour for sucrose would eventually reduce in intensity at a decreasing rate until it becomes constant.

- This is because sucrose would continue to decrease in the medium until it is depleted
- Also, the accumulation of alcohol in the medium would eventually inhibit the action of the enzymes of the yeast cells.

(e) (i) – to prevent the escape of the carbon dioxide produced during process of yeast respiration.

- To prevent entry of oxygen into the reaction medium. This would lead to rapid aerobic breakdown of the food substrates.

(ii) – to maintain the temperature at a value optimum for the functioning of the yeast enzymes. This is important in order to ensure that only the investigated variables are investigated.