

UCE

Mathematics 3

(For S.3)

BY

Chepa Y. M

Department of Mathematics, Kibuli Secondary School.
Telephone: +256 772 481628 Email: ymussy@yahoo.com



Second edition 2008

© Chepa Y. M. March 2008

All Rights Reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the author.



Contents

Chapter	Page
1. Vector Geometry	4
Vector addition and subtraction.....	4
Multiplication by a scalar	5
Parallel vectors.....	5
Collinearity	6
2. Matrices	16
Elementary operations with matrices.....	16
The Determinant of a 2×2 Matrix	24
The inverse of a 2×2 matrix	24
3 Matrices and Transformations	29
Translations	29
Rotations	30
Reflections.....	31
Enlargements.....	32
Shears.....	33
Stretches	34
Transformation matrices	35
Successive transformations.....	42
Matrices of successive transformations	46
The inverse of a transformation	49
Area and the determinant of a matrix.....	51
4. Statistics	57
Methods of presentation of data.....	58
Pie Chart.....	58
Frequency Distributions	59
Cumulative frequency	61
Histograms.....	62
Frequency polygons.	64
Cumulative frequency curve (Ogive)	66
The mean	67
The Median.....	70
The mode	73
Quartiles	75
5. Probability	84
Definitions.....	84

Review of terms used in set theory	84
Independent events	85
Venn diagrams	85
Basic rules of probability	86
Tree diagrams	90
6. Sets.....	97
Symbols, Problems involving three sets	97
7. Formulae.....	105
subject of the formula.....	105
Substitution	109
8. Functions.....	112
Inverse of a function	115
Composite functions	116
Rational functions	118
9. Trigonometry (2).....	120
Tan, sin and cos of 45°	120
Tan, sin and cos of 60° and 30°	121
The graph of $y = \sin x$ (Sine Curve)	122
The graph of $y = \cos x$	123
The sine rule	123
The Cosine Rule.....	126
The area of a triangle.....	127
10. Three dimensional geometry	132
Geometrical properties of solids.....	133
Projection of a line on a plane.....	134
An angle between two lines	135
Skew lines	137
The length of a line on a solid	140
Angle between a line and a plane	143
Angle between two planes.....	147
11. Further Logarithms.....	158
Logarithmic notation.....	158
Laws of logarithms.....	158
Simplification of logarithmic expressions	159
Solution of logarithmic equations.....	162

1. Vector Geometry

1.1 Revision of Vectors

A scalar is a quantity with magnitude (size) only.

A vector is a quantity with both magnitude and direction.


Velocity is speed in a given direction and is thus a vector.

e.g 15 m/s is a speed.

15 m/s on a bearing 070° is a velocity since the direction is given.

A vector can be represented on paper by a line segment, say AB, showing magnitude and direction or by an ordered set of numbers. We can write vector

$$\mathbf{AB} \text{ as } \mathbf{AB} = \mathbf{a} = \begin{pmatrix} 4 \\ 3 \end{pmatrix}$$



When writing these vectors you will use the common notation \vec{AB} or \underline{a}

$\mathbf{AB} = \begin{pmatrix} 4 \\ 3 \end{pmatrix}$ is referred to as **column** or **displacement** vector.

A **position vector** starts from the origin. If P is the point (x,y) and O is the origin,

then the position vector of P is $\mathbf{OP} = \begin{pmatrix} x \\ y \end{pmatrix}$

Also, if A and B have position vectors \mathbf{a} and \mathbf{b} with reference to the origin O, then $\mathbf{AB} = \mathbf{b} - \mathbf{a}$.

Vector addition and subtraction

- (i) Vectors are added by the Triangle Rule of vector addition, i.e. where vector \mathbf{a} ends \mathbf{b} starts.

$$\mathbf{AB} + \mathbf{BC} = \mathbf{AC}$$



- (ii) The triangular rule follows from the parallelogram law. If \mathbf{AB} and \mathbf{AD} are two non-parallel vectors drawn through A, then the sum $\mathbf{AB} + \mathbf{AD}$ is called the resultant vector.

The resultant is \mathbf{AC} , the diagonal through A of the parallelogram whose sides are AB and AD.

$$\mathbf{AB} + \mathbf{AD} = \mathbf{AB} + \mathbf{BC} = \mathbf{AC} = \mathbf{a} + \mathbf{b}$$

$$\mathbf{AD} + \mathbf{DC} = \mathbf{AC} = \mathbf{b} + \mathbf{a}.$$

$$\text{Hence } \mathbf{a} + \mathbf{b} = \mathbf{b} + \mathbf{a}$$




Subtraction can also be performed in a similar way.

$\mathbf{CB} = -\mathbf{BC}$. (The negative sign reverses the direction of the vector).

$$\mathbf{AB} - \mathbf{CB} = \mathbf{AB} - (-\mathbf{BC})$$

$$= \mathbf{AB} + \mathbf{BC}$$

$$= \mathbf{AC}.$$



Multiplication by a scalar

When a vector $\begin{pmatrix} a \\ b \end{pmatrix}$ is multiplied by a scalar k , we get $k \begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} ka \\ kb \end{pmatrix}$. The scalar k

can be any positive or negative number. Each component of the vector is multiplied by the scalar.

Parallel vectors

If $\mathbf{a} = k\mathbf{b}$, where k is a scalar, then \mathbf{a} is parallel to \mathbf{b} . Conversely, if \mathbf{a} is parallel to \mathbf{b} , then $\mathbf{a} = k\mathbf{b}$.

For example, $\begin{pmatrix} 3 \\ -5 \end{pmatrix}$ is parallel to $\begin{pmatrix} 6 \\ -10 \end{pmatrix}$, because $\begin{pmatrix} 6 \\ -10 \end{pmatrix}$ may be written

$$2 \begin{pmatrix} 3 \\ -5 \end{pmatrix}.$$


In general the vector $k \begin{pmatrix} a \\ b \end{pmatrix}$ is parallel to $\begin{pmatrix} a \\ b \end{pmatrix}$.

Collinearity

If $\mathbf{AB} = k\mathbf{BC}$, where k is a scalar, then \mathbf{AB} is parallel to \mathbf{BC} .
But B is common, so A , B and C lie on a straight line, i.e. the points A , B and C are **collinear**.

Proportional division of a line

(a)




$AP : PB = 3 : 1$, then we say that P divides AB internally in the ratio $3 : 1$.
(P lies between A and B).

Note: The direction is important as P divides BA in the ratio $1 : 3$.

$$\mathbf{AP} = \frac{3}{4}\mathbf{AB} \text{ and } \mathbf{PB} = \frac{1}{4}\mathbf{AB}$$

(b)




$$\mathbf{AQ} : \mathbf{QB} = 4 : -1.$$

The negative sign is required as QB is in the opposite direction to AB . We say that Q divides AB externally in the ratio $4 : 1$, or Q divides AB in the ratio $4 : -1$.

Whenever Q is outside AB (on either side), $\mathbf{AQ} : \mathbf{QB}$ will be negative, and we say that Q divides AB externally.

Example 1.1

Write down the relationship between the vectors \mathbf{a} , \mathbf{b} and \mathbf{c} .




Solution

Using the vector law of addition, $\underline{\mathbf{a}} + \underline{\mathbf{b}} = \underline{\mathbf{c}}$

Example 1.2

In $\triangle OCD$, $OC = c$ and $OD = d$.




- Express CD in terms of vectors c and d .
- M is the midpoint of CD . What is CM in terms of c and d ?
- Using your answers to (a) and (b), find OM in terms of c and d .

Solution

- $CD = CO + OD = -OC + OD$
 $= -c + d$ or $d - c$
Therefore, $CD = d - c$.
- $CM = \frac{1}{2} CD$
 $= \frac{1}{2} (d - c)$.
- $OM = OC + CM$
 $= c + \frac{1}{2} (d - c) = c - \frac{1}{2} c + \frac{1}{2} d$
 $= \frac{1}{2} c + \frac{1}{2} d$
 $= \frac{1}{2} (c + d)$

Example 1.3

In the diagram below, P divides the line AB in the ratio $AP : PB = 7 : 3$. If $OA = a$ and $OB = b$, express OP in terms of a and b .



Solution


- In triangle OAB , $OA + AB = OB$
 $a + AB = b$
 $AB = b - a$

Along AB, $\overrightarrow{AP} = \frac{7}{10}\overrightarrow{AB} = \frac{7}{10}(\mathbf{b}-\mathbf{a})$

$$\begin{aligned}\overrightarrow{OP} &= \overrightarrow{OA} + \overrightarrow{AP} = \mathbf{a} + \frac{7}{10}(\mathbf{b}-\mathbf{a}) \\ &= \frac{3}{10}\mathbf{a} + \frac{7}{10}\mathbf{b}\end{aligned}$$

Example 1.4

In the figure below, $\overrightarrow{OA} = \mathbf{a}$ and $\overrightarrow{OB} = \mathbf{b}$.



- Express \overrightarrow{BA} in terms of \mathbf{a} and \mathbf{b} .
- If X is the mid-point of \overrightarrow{BA} , show that $\overrightarrow{OX} = \frac{1}{2}(\mathbf{a} + \mathbf{b})$.
- Given that $\overrightarrow{OC} = 3\mathbf{a}$, express \overrightarrow{BC} in terms of \mathbf{a} and \mathbf{b} .
- Given that $\overrightarrow{BY} = m\overrightarrow{BC}$, express \overrightarrow{OY} in terms of \mathbf{a} , \mathbf{b} , m .
- If $\overrightarrow{OY} = n\overrightarrow{OX}$ use the results of (b) and (d) to evaluate m and n .

Solution

- In triangle OAP,
 $\overrightarrow{OB} + \overrightarrow{BA} = \overrightarrow{OA}$
 $\mathbf{b} + \overrightarrow{BA} = \mathbf{a}$
 $\overrightarrow{BA} = \mathbf{a} - \mathbf{b}$
- In triangle OBX, $\overrightarrow{BX} = \frac{1}{2}\overrightarrow{BA}$
 $= \frac{1}{2}(\mathbf{a} - \mathbf{b})$
 $\overrightarrow{OX} = \overrightarrow{OB} + \overrightarrow{BX} = \mathbf{b} + \frac{1}{2}(\mathbf{a} - \mathbf{b})$
 $= \frac{1}{2}(\mathbf{a} + \mathbf{b})$
- In $\triangle OBC$, $\overrightarrow{BC} = \overrightarrow{BO} + \overrightarrow{OC}$
 $= -\mathbf{b} + 3\mathbf{a} = 3\mathbf{a} - \mathbf{b}$
- In $\triangle OBY$, $\overrightarrow{OY} = \overrightarrow{OB} + \overrightarrow{BY}$
 $= \mathbf{b} + m(3\mathbf{a} - \mathbf{b})$
 $= 3m\mathbf{a} + (1 - m)\mathbf{b}$ (i)
- $\overrightarrow{OY} = n\overrightarrow{OX} = n\frac{1}{2}(\mathbf{a} + \mathbf{b}) = \frac{1}{2}n\mathbf{a} + \frac{1}{2}n\mathbf{b}$ (ii)
 Also, $\overrightarrow{OY} = 3m\mathbf{a} + (1 - m)\mathbf{b}$

Since the vectors in (i) and (ii) are identical, the scalars multiplying \mathbf{a} and \mathbf{b} can be equated:

$$\frac{1}{2}n = 3m \quad (\text{scalars of } \mathbf{a})$$

$$\frac{1}{2}m = 1 - m \quad (\text{scalars of } \mathbf{b})$$

$$3m = 1 - m \Leftrightarrow 4m = 1 \Leftrightarrow m = \frac{1}{4}$$

$$\text{If } m = \frac{1}{4}, \text{ then } \frac{1}{2}n = 3 \times \frac{1}{4} \Leftrightarrow n = 1\frac{1}{2}$$

Notice the last stage of example 1.4. In general if


$$h\mathbf{a} + k\mathbf{b} = n\mathbf{a} + m\mathbf{b}, \text{ then}$$

$$h = n \text{ and } k = m \text{ (or } \mathbf{a} = \mathbf{0} \text{ and } \mathbf{b} = \mathbf{0})$$

Example 1.5

In the diagram, $OD = 2OA$, $OE = 4OB$, $OA = \mathbf{a}$ and $OB = \mathbf{b}$.

- Express the following vectors in terms of \mathbf{a} and \mathbf{b} .
 OD , OE , BA and ED .
- Given that $BC = 3BA$, express:
 - OC
 - EC , in terms of \mathbf{a} and \mathbf{b} .
- Hence show that the points E , D and C lie on a straight line.



Solution

$$(a) \quad OD = 2OA = 2\mathbf{a};$$

$$OE = 4OB = 4\mathbf{b}$$

$$\begin{aligned} BA &= BO + OA = -OB + OA \\ &= OA - OB \\ &= \mathbf{a} - \mathbf{b} \end{aligned}$$

$$\begin{aligned} ED &= EO + OD = -OE + OD \\ &= OD - OE \\ &= 2\mathbf{a} - 4\mathbf{b} \text{ or } 2(\mathbf{a} - 2\mathbf{b}) \end{aligned}$$

$$(b) \quad (i) \quad OC = OB + BC = \mathbf{b} + 3BA$$

$$\begin{aligned}
&= \mathbf{b} + 3(\mathbf{a} - \mathbf{b}) \\
&= \mathbf{b} - 3\mathbf{b} + 3\mathbf{a} \\
&= -2\mathbf{b} + 3\mathbf{a} \text{ or } 3\mathbf{a} - 2\mathbf{b}
\end{aligned}$$

$$\therefore \mathbf{OC} = 3\mathbf{a} - 2\mathbf{b}.$$


$$\begin{aligned}
\text{(ii) } \mathbf{EC} &= \mathbf{EO} + \mathbf{OC} = -4\mathbf{b} + (3\mathbf{a} - 2\mathbf{b}) \\
&= -4\mathbf{b} - 2\mathbf{b} + 3\mathbf{a} \\
&= -6\mathbf{b} + 3\mathbf{a} \text{ or } 3(\mathbf{a} - 2\mathbf{b})
\end{aligned}$$

$$\therefore \mathbf{EC} = 3(\mathbf{a} - 2\mathbf{b})$$

- (c) Using results for \mathbf{ED} and \mathbf{DC} , we see that $\mathbf{EC} = \frac{3}{2}\mathbf{ED}$. This means that \mathbf{EC} and \mathbf{ED} are parallel vectors. Since both vectors pass through the point E, the points E, D and C must lie on a straight line.

Example 1.6

In the figure below, OABC is a parallelogram, M is the mid-point of OA and $\mathbf{AX} = \frac{2}{7}\mathbf{AC}$. $\mathbf{OA} = \mathbf{a}$ and $\mathbf{OC} = \mathbf{c}$.



- (a) Express the following in terms of \mathbf{a} and \mathbf{c} .
- (i) \mathbf{MA} (ii) \mathbf{AB} (iii) \mathbf{AC} (iv) \mathbf{AX}
- (b) Using triangle MAX, express \mathbf{MX} in terms of \mathbf{a} and \mathbf{c} .
- (c) If $\mathbf{AY} = p\mathbf{AB}$, use triangle MAY to express \mathbf{MY} in terms of \mathbf{a} , \mathbf{c} and p .
- (d) Also if $\mathbf{MY} = q\mathbf{MX}$, use the result in (b) to express \mathbf{MY} in terms of \mathbf{a} , \mathbf{c} and q .
- (e) Hence find p and q and the ratio $\mathbf{AY} : \mathbf{YB}$

Solution

- (a) (i) $\mathbf{MA} = \frac{1}{2}\mathbf{OA} = \frac{1}{2}\mathbf{a}$.
- (ii) $\mathbf{AB} = \mathbf{OC}$ since OABC is a parallelogram.
 $\therefore \mathbf{AB} = \mathbf{c}$.
- (iii) $\mathbf{AC} = \mathbf{AO} + \mathbf{OC}$
 $= -\mathbf{a} + \mathbf{c}$ or $\mathbf{c} - \mathbf{a}$
- (iv) $\mathbf{AX} = \frac{2}{7}\mathbf{AC} = \frac{2}{7}(\mathbf{c} - \mathbf{a})$
- (b) $\mathbf{MX} = \mathbf{MA} + \mathbf{AX}$

$$= \frac{1}{2}\mathbf{a} + \frac{2}{7}(\mathbf{c} - \mathbf{a}) = \frac{7\mathbf{a} + 4\mathbf{c} - 4\mathbf{a}}{14}$$

$$= \frac{3\mathbf{a} + 4\mathbf{c}}{14} \text{ or } \frac{3}{14}(\mathbf{a} + \mathbf{c})$$

$$\therefore \mathbf{MX} = \frac{3}{14}(\mathbf{a} + \mathbf{c}).$$

(c) $\mathbf{MY} = \mathbf{MA} + \mathbf{AY}$
 $= \frac{1}{2}\mathbf{a} + p\mathbf{AB} = \frac{1}{2}\mathbf{a} + p\mathbf{c} \text{ or } \frac{1}{2}(\mathbf{a} + 2p\mathbf{c})$

(d) If $\mathbf{MY} = q\mathbf{MX}$, then $\mathbf{MY} = q\frac{3}{14}(\mathbf{a} + \mathbf{c})$.

(e) From (c) and (d), we equate the expressions for \mathbf{MY} :

$$\frac{1}{2}(\mathbf{a} + 2p\mathbf{c}) = q\frac{3}{14}(\mathbf{a} + \mathbf{c}).$$

Equating the scalars multiplying \mathbf{a} and \mathbf{c} gives,

For \mathbf{a} : $\frac{1}{2} = \frac{3}{14}q$ (i)

For \mathbf{c} : $p = \frac{3}{14}q$ (ii)

From (i), $q = \frac{7}{3}$.

Substituting for q into (ii) gives,

$$p = \frac{7}{3} \times \frac{3}{14} = \frac{1}{2}$$


Hence, $\mathbf{AY} = \frac{1}{2}\mathbf{AB}$ and $\mathbf{AY} : \mathbf{YB} = 1:1$

Exercise 1.1


Make sketches where necessary.

1. Using the figure below represent each of the following by a single vector.

- | | |
|---|---|
| (a) $\mathbf{PQ} + \mathbf{QR}$ | (b) $\mathbf{PR} + \mathbf{RS}$ |
| (c) $\mathbf{PS} + \mathbf{ST}$ | (d) $\mathbf{PR} + \mathbf{RT}$ |
| (e) $\mathbf{PQ} + \mathbf{QR} + \mathbf{RS}$ | (f) $\mathbf{PQ} + \mathbf{QT} + \mathbf{TS}$ |
| (g) $\mathbf{PQ} + \mathbf{QR} + \mathbf{RS}$ | (h) $\mathbf{PQ} + \mathbf{QT} + \mathbf{TR} + \mathbf{RS}$ |




2. In the diagram, $OA = a$, $OB = b$ and M is the mid-point of AB .




Find OM in terms of a and b .

3. Express XY , YZ and ZX in terms of a , b and c .




4. $OP = 2a$, $PQ = 2b - 3a$, $OR = 3b$.




Express (a) OQ , (b) QR in terms of a and b as simply as possible.

5. $ORST$ is a parallelogram, $OR = r$ and $OT = t$
If $ST = 4SP$, express the following in terms of r and or t .

- | | |
|----------|----------|
| (a) RS | (b) ST |
| (c) SP | (d) RP |
| (e) QP | |



6. In $\triangle PQR$, $PQ = \mathbf{a}$, $PR = \mathbf{b}$ and S is the mid-point of PR. Express the following in terms of \mathbf{a} and or \mathbf{b} .
- (a) QR (b) PS
 (c) QS
7. In $\triangle PQR$, A is a point on PR such that $PA = \frac{4}{5} PR$ and B is the mid-point of QR. Point C lies on PQ produced so that $PC = \frac{3}{2} PQ$. If $PR = x$ and $PQ = y$ express the following in terms of x and y .
- (a) PA (b) PB
 (c) PC (d) AB
8. In the diagram, P is a point on AB such that $BA = 4BP$ and Q is the mid-point of OA. OP and BQ intersect at X.




Given $OA = \mathbf{a}$ and $OB = \mathbf{b}$:


- (a) Express the following in terms of \mathbf{a} and \mathbf{b} .
- (i) AB (ii) OP
 (iii) BQ
- (b) If $BX = hBQ$, express OX in terms of \mathbf{a} , \mathbf{b} and h .
- (c) If $OX = kOP$ use the previous result to find h and k .
- (d) Hence express OX in terms of \mathbf{a} and \mathbf{b} only.
9. (a) Given that: $OY = 2OB$, $OX = \frac{5}{2} OA$, $OA = \mathbf{a}$ and $OB = \mathbf{b}$, express the following in terms of \mathbf{a} and \mathbf{b} .

OY, OX, AB, and XY.


- (b) Given that $AC = 6AB$, express OC and XC in terms of \mathbf{a} and \mathbf{b} .
 (c) Use the results for XY and XC to show that points X, Y and C lie on a straight line.




10. In the diagram, $OA = \mathbf{a}$ and $OB = \mathbf{b}$, M is the mid-point of OA and P lies on AB such that $AP = \frac{2}{3}AB$.
- (a) Express the following in terms of \mathbf{a} and \mathbf{b} : AB , AP , MA and MP
 (b) If X lies on OB produced such that $OB = BX$, express MX in terms of \mathbf{a} and \mathbf{b} .
 (c) Show that MPX is a straight line.



11. X is the mid-point of OD, Y lies on CD such that $CY = \frac{1}{4}CD$, $OC = \mathbf{c}$ and $OD = \mathbf{d}$.
- (a) Express the following in terms of \mathbf{c} and \mathbf{d} : CD , CY , CX and OY .
 (b) Given that $CZ = hCX$, express OZ in terms of \mathbf{c} , \mathbf{d} and h .
 (c) If $OZ = kOY$, form an equation and hence find the values of h and k .



12. In the diagram $OA = \mathbf{a}$ and $OB = \mathbf{b}$. The points P and Q are such that $AP = \frac{1}{3}AB$ and $OQ = \frac{1}{3}OA$. Express OP and BQ in terms of \mathbf{a} and \mathbf{b} . If $OC = h OP$ and $BC = k BQ$, express OC in two different ways and hence deduce
- the values of h and k ,
 - the vector OC in terms of \mathbf{a} and \mathbf{b} only
 - the ratio in which C divides BQ .



13. In $\triangle OXY$, $OX = \mathbf{x}$ and $OY = \mathbf{y}$. The point A lies on OY such that $OA : AY = 1 : 3$. The point B lies on XY such that $XB : BY = 2 : 3$. Find in terms of \mathbf{x} and \mathbf{y} the vectors OA , XY , XB , OB and AX . The lines OB and AX intersect at point C. Given that $OC = h OB$ and $AC = k AX$, express the position vector of C in terms of:
- h , \mathbf{x} and \mathbf{y} ,
 - k , \mathbf{x} and \mathbf{y} .
- Hence find the values of h and k and determine the ratios $OC : CB$ and $AC : CX$.
14. In $\triangle OPQ$; $OP = \mathbf{p}$, $OQ = \mathbf{q}$. A and B are points on OQ and OP respectively, such that $OA : AQ = 3 : 1$ and $OB = \frac{1}{4}OP$. AP and BQ intersect at C.
- Show that $OC = \frac{1}{13}(\mathbf{p} + 9\mathbf{q})$.
 - Determine the ratios $AC : CP$ and $BC : CQ$.

2. Matrices

A matrix is a rectangular array of numbers. It is enclosed by large curved brackets. Each number of the array is called an entry (or an element). The elements are arranged in rows and columns. Rows are across the page, i.e. horizontal, and columns are vertical.

Matrices (plural of matrix) are often used for data storage.

Matrices are represented by capital letters and written with a wavy line underneath. For example, a matrix can be written as

$$\mathbf{A} = \underline{\underline{A}} = \begin{pmatrix} 2 & 4 & -3 \\ 1 & 7 & 0 \end{pmatrix}. \text{ In text books it is printed in bold } \mathbf{A}.$$

Order of a matrix

The order of a matrix is the number of rows and columns in the matrix. For example, $\begin{pmatrix} 2 & 4 & -3 \\ 1 & 7 & 0 \end{pmatrix}$ is of order 2×3 (read as 'two by three') because it has 2

rows and 3 columns. Notice that the number of rows is given first. This can be remembered by the first letters of the words **R**esistance **C**ouncil: $\mathbf{R} \times \mathbf{C}$.

If the number of rows in a matrix equals the number of columns, the matrix is called a **square matrix**. For example,

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

A matrix with only one row is called a **row matrix**.

A matrix with only one column is called a **column matrix**.

Elementary operations with matrices

Two matrices of the same order can be added or subtracted by adding (or subtracting) the corresponding elements of the matrices to give a third matrix.

$$\text{For example, } \begin{pmatrix} -2 & 1 \\ 3 & 7 \\ 6 & 9 \end{pmatrix} + \begin{pmatrix} 1 & 1 \\ -1 & 0 \\ 3 & 1 \end{pmatrix} = \begin{pmatrix} -2+1 & 1+1 \\ 3-1 & 7+0 \\ 6+3 & 9+1 \end{pmatrix} = \begin{pmatrix} -1 & 2 \\ 2 & 7 \\ 9 & 10 \end{pmatrix}$$

$$\text{And } \begin{pmatrix} -2 & 1 \\ 3 & 7 \\ 6 & 9 \end{pmatrix} - \begin{pmatrix} 1 & 1 \\ -1 & 0 \\ 3 & 1 \end{pmatrix} = \begin{pmatrix} -2-1 & 1-1 \\ 3-(-1) & 7-0 \\ 6-3 & 9-1 \end{pmatrix} = \begin{pmatrix} -3 & 0 \\ 4 & 7 \\ 3 & 8 \end{pmatrix}.$$

Note: Given any two matrices \mathbf{A} and \mathbf{B} , then $\mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A}$ (commutative property).

Also, $(\mathbf{A} + \mathbf{B}) + \mathbf{C} = \mathbf{A} + (\mathbf{B} + \mathbf{C})$ (Associative property).

Multiplication of a matrix by a scalar

To multiply a matrix \mathbf{A} by a scalar k (a real number), we multiply each number,



i.e. each element of the matrix by k.

For example, if $P = \begin{pmatrix} 1 & 1 & 7 \\ 0 & 2 & 6 \end{pmatrix}$ then $3P = 3 \times \begin{pmatrix} 1 & 1 & 7 \\ 0 & 2 & 6 \end{pmatrix} = \begin{pmatrix} 3 & 3 & 21 \\ 0 & 6 & 18 \end{pmatrix}$.

Matrix multiplication

In matrix multiplication, each row of the first matrix is *multiplied* by each column of the second matrix. In the multiplication of a row by a column, each element in the row is *multiplied* by the corresponding element in the column and then the

products *added*. For example, $(a \ b \ c) \begin{pmatrix} p \\ q \\ r \end{pmatrix} = (ap + bq + cr)$.

Note: Two matrices can be multiplied if the number of columns in the first matrix is equal to the number of rows in the second matrix. For example, if

$P = \begin{pmatrix} 3 & 7 \\ 2 & 6 \\ 1 & 0 \end{pmatrix}$ and $Q = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$ then PQ can be found as P has 2 columns and Q has

2 rows.

$$PQ = \begin{pmatrix} 3 & 7 \\ 2 & 6 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix} = \begin{pmatrix} 3 \times 1 + 7 \times 3 & 3 \times 2 + 7 \times 4 \\ 2 \times 1 + 6 \times 3 & 2 \times 2 + 6 \times 4 \\ 1 \times 1 + 0 \times 3 & 1 \times 2 + 0 \times 4 \end{pmatrix} = \begin{pmatrix} 3+21 & 6+28 \\ 2+18 & 4+24 \\ 1+0 & 2+0 \end{pmatrix} = \begin{pmatrix} 24 & 34 \\ 20 & 28 \\ 1 & 2 \end{pmatrix}$$

P has order 3×2 , Q has order 2×2 and so PQ has order 3×2 .

QP cannot be found because the number of rows in Q is not equal to the number of columns in P .

Note:

- (i) In general $PQ \neq QP$. (Matrix multiplication is not commutative).
- (ii) If P is $(m \times n)$ matrix and Q is $(n \times p)$ matrix, then Q can be premultiplied by P .

$$\begin{matrix} P & \times & Q & = & R \\ (m \times n) & & (n \times p) & & (m \times p) \end{matrix}$$

- (iii) In matrices A^2 means $A \times A$. You must multiply the matrices together. Whereas, $2A$ means $2 \times A$.

Example 2.1

Perform the following multiplications.

(a) $\begin{pmatrix} 3 & 2 \\ 4 & 1 \end{pmatrix} \begin{pmatrix} 2 & 1 \\ 1 & 5 \end{pmatrix}$ (b) $\begin{pmatrix} 2 & 1 & -2 \\ 0 & 1 & 3 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & -2 \\ 4 & 3 \end{pmatrix}$

Solution

(a) $\begin{pmatrix} 3 & 2 \\ 4 & 1 \end{pmatrix} \begin{pmatrix} 2 & 1 \\ 1 & 5 \end{pmatrix} = \begin{pmatrix} 6+2 & 3+10 \\ 8+1 & 4+5 \end{pmatrix} = \begin{pmatrix} 8 & 13 \\ 9 & 9 \end{pmatrix}$



$$(b) \quad \begin{pmatrix} 2 & 1 & -2 \\ 0 & 1 & 3 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & -2 \\ 4 & 3 \end{pmatrix} = \begin{pmatrix} 2+1-8 & 0-2-6 \\ 0+1+12 & 0-2+9 \end{pmatrix} \\ = \begin{pmatrix} -5 & -8 \\ 13 & 7 \end{pmatrix}$$

Example 2.2

If $A = \begin{pmatrix} 1 & 0 \\ 3 & 2 \end{pmatrix}$, $B = \begin{pmatrix} x & 0 \\ 1 & 3 \end{pmatrix}$, and $AB = BA$, find x .

Solution

$$AB = \begin{pmatrix} 1 & 0 \\ 3 & 2 \end{pmatrix} \begin{pmatrix} x & 0 \\ 1 & 3 \end{pmatrix} = \begin{pmatrix} x+0 & 0+0 \\ 3x+2 & 0+6 \end{pmatrix} = \begin{pmatrix} x & 0 \\ 3x+2 & 6 \end{pmatrix}$$

$$BA = \begin{pmatrix} x & 0 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 3 & 2 \end{pmatrix} = \begin{pmatrix} x+0 & 0+0 \\ 1+9 & 0+6 \end{pmatrix} = \begin{pmatrix} x & 0 \\ 10 & 6 \end{pmatrix}$$

Equating AB and BA , we get

$$\begin{pmatrix} x & 0 \\ 3x+2 & 6 \end{pmatrix} = \begin{pmatrix} x & 0 \\ 10 & 6 \end{pmatrix}$$

Note: When two matrices are equal, then their corresponding elements must also be equal. Thus,

$$3x + 2 = 10$$

$$\Rightarrow 3x = 8$$

$$\Rightarrow x = \frac{8}{3}$$

Example 2.3

If $\begin{pmatrix} 2x & 1 \\ 0 & 4 \end{pmatrix} + \begin{pmatrix} 3 & 2 \\ 2 & 1 \end{pmatrix} = \begin{pmatrix} 9 & 3 \\ 2 & 5 \end{pmatrix}$, find the value of x .

Solution

$$\begin{pmatrix} 2x & 1 \\ 0 & 4 \end{pmatrix} + \begin{pmatrix} 3 & 2 \\ 2 & 1 \end{pmatrix} = \begin{pmatrix} 2x+3 & 3 \\ 2 & 5 \end{pmatrix} = \begin{pmatrix} 9 & 3 \\ 2 & 5 \end{pmatrix}$$

By equating the corresponding elements of both sides, we get

$$2x + 3 = 9 \Rightarrow 2x = 6 \Rightarrow x = 3.$$

Example 2.4

Given that $B = \begin{pmatrix} 5 & 5 \\ -2 & -2 \end{pmatrix}$, find k if $B^2 = kB$.

Solution

$$\mathbf{B}^2 = \begin{pmatrix} 5 & 5 \\ -2 & -2 \end{pmatrix} \begin{pmatrix} 5 & 5 \\ -2 & -2 \end{pmatrix} = \begin{pmatrix} 25-10 & 25-10 \\ -10+4 & -10+4 \end{pmatrix} = \begin{pmatrix} 15 & 15 \\ -6 & -6 \end{pmatrix}$$

It can be seen that $\mathbf{B}^2 = \begin{pmatrix} 15 & 15 \\ -6 & -6 \end{pmatrix} = 3 \times \begin{pmatrix} 5 & 5 \\ -2 & -2 \end{pmatrix} = 3\mathbf{B}$

Therefore, the value of k for which $\mathbf{B}^2 = k\mathbf{B}$ is 3.

Example 2.5

Given $\mathbf{P} = \begin{pmatrix} 2 & -3 \\ 2 & 1 \end{pmatrix}$, $\mathbf{Q} = \begin{pmatrix} 14 & 2 \\ -8 & 6 \end{pmatrix}$ and $\mathbf{R} = \begin{pmatrix} 16 & 20 \\ 32 & 4 \end{pmatrix}$ calculate:

(a) $5\mathbf{Q} - 3\mathbf{P}$

(b) $\frac{1}{2}\mathbf{Q} + \mathbf{R}$

Solution

$$\begin{aligned} \text{(a) } 5\mathbf{Q} - 3\mathbf{P} &= 5 \begin{pmatrix} 14 & 2 \\ -8 & 6 \end{pmatrix} - 3 \begin{pmatrix} 2 & -3 \\ 2 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 70 & 10 \\ -40 & 30 \end{pmatrix} - \begin{pmatrix} 6 & -9 \\ 6 & 3 \end{pmatrix} \\ &= \begin{pmatrix} 70-6 & 10+9 \\ -40-6 & 30-3 \end{pmatrix} = \begin{pmatrix} 64 & 19 \\ -46 & 27 \end{pmatrix} \end{aligned}$$

$$\begin{aligned} \text{(b) } \frac{1}{2}\mathbf{Q} + \mathbf{R} &= \frac{1}{2} \times \begin{pmatrix} 14 & 2 \\ -8 & 6 \end{pmatrix} + \begin{pmatrix} 16 & 20 \\ 32 & 4 \end{pmatrix} \\ &= \begin{pmatrix} 7 & 1 \\ -4 & 3 \end{pmatrix} + \begin{pmatrix} 16 & 20 \\ 32 & 4 \end{pmatrix} = \begin{pmatrix} 7+16 & 1+20 \\ -4+32 & 3+4 \end{pmatrix} \\ &= \begin{pmatrix} 23 & 21 \\ 28 & 7 \end{pmatrix}. \end{aligned}$$

Example 2.6

Three garages G_1 , G_2 and G_3 sell cars of two types A and B. The sales in one week are shown in the table below together with a matrix showing the price paid for each type.

Sales:

Garage	Type	
	A	B
G_1	3	1
G_2	2	0
G_3	4	1

Prices (in dollars): $\begin{pmatrix} \text{A: } 5000 \\ \text{B: } 7500 \end{pmatrix}$

Write down the sales as a 3×2 matrix and hence work out the total value of sales for the three garages.

Solution

Sales:

$$\begin{array}{l} \mathbf{G}_1 \\ \mathbf{G}_2 \\ \mathbf{G}_3 \end{array} \begin{array}{cc} \mathbf{A} & \mathbf{B} \\ \left(\begin{array}{cc} 3 & 1 \\ 2 & 0 \\ 4 & 1 \end{array} \right) \end{array}$$

The total value of sales for the three garages is given by the matrix product

$$\begin{pmatrix} 3 & 1 \\ 2 & 0 \\ 4 & 1 \end{pmatrix} \begin{pmatrix} 5000 \\ 7500 \end{pmatrix} = \begin{pmatrix} 15000 + 7500 \\ 10000 + 0 \\ 20000 + 7500 \end{pmatrix} = \begin{pmatrix} 22500 \\ 10000 \\ 27500 \end{pmatrix}$$

The total sales for each garage are therefore given in the following matrix

$$\begin{array}{l} \mathbf{G}_1 \\ \mathbf{G}_2 \\ \mathbf{G}_3 \end{array} \begin{array}{c} \text{Value of sales (\$)} \\ \left(\begin{array}{c} 22500 \\ 10000 \\ 27500 \end{array} \right) \end{array}$$

Zero (or null) matrix

A matrix with all its elements zero is called a **zero** matrix. If to a given matrix **A**, a zero matrix (of the same order) is added, the value of **A** is unchanged. However, if we multiply a matrix by the zero matrix we get a zero matrix. Thus, $\mathbf{A} + \mathbf{O} = \mathbf{A}$ and $\mathbf{AO} = \mathbf{OA} = \mathbf{O}$, where **O** is the zero matrix.

Identity (or Unit Matrix)

A square matrix in which the elements of the leading (or major) diagonal are 1 and the other elements are zero is called the **identity matrix**, and is denoted by **I**.

Thus, an identity matrix of order 2×2 will be $\mathbf{I} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$.

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

← Leading or major diagonal.

An identity matrix of order 3 will be

$$\mathbf{I} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

If $\mathbf{A} = \begin{pmatrix} 2 & 3 \\ 1 & 4 \end{pmatrix}$ is a 2×2 matrix, then $\mathbf{AI} = \begin{pmatrix} 2 & 3 \\ 1 & 4 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 2 & 3 \\ 1 & 4 \end{pmatrix} = \mathbf{A}$

$$\text{And } \mathbf{IA} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 2 & 3 \\ 1 & 4 \end{pmatrix} = \begin{pmatrix} 2 & 3 \\ 1 & 4 \end{pmatrix} = \mathbf{A}.$$

Hence, $\mathbf{AI} = \mathbf{IA} = \mathbf{A}$.

$$\text{Also } \mathbf{I}^2 = \mathbf{I} \times \mathbf{I} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} = \mathbf{I}$$

In general, $\mathbf{I}^n = \mathbf{I}$.

Exercise 2.1

In questions 1 - 3, find the value of the following

$$1. \quad \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix} + \begin{pmatrix} 2 \\ 1 \\ 3 \end{pmatrix} \qquad 2. \quad 2 \begin{pmatrix} 1 \\ 3 \\ 1 \\ 5 \end{pmatrix} + 4 \begin{pmatrix} 1 \\ 0 \\ 1 \\ 6 \end{pmatrix}$$

$$3. \quad 8 \begin{pmatrix} 1 & 2 \\ 5 & 3 \end{pmatrix} - 2 \begin{pmatrix} 2 & 13 \\ 17 & 16 \end{pmatrix}$$

$$4. \quad \text{If } \mathbf{A} = \begin{pmatrix} 1 & 2 \\ 3 & 1 \end{pmatrix}, \mathbf{B} = \begin{pmatrix} 5 & -3 \\ 6 & 2 \end{pmatrix} \text{ and } \mathbf{C} = \begin{pmatrix} 5 & 3 \\ -4 & 2 \end{pmatrix}$$

(a) Calculate $\mathbf{A} + \mathbf{B}$ and $\mathbf{B} + \mathbf{A}$.

(b) Calculate: (i) $3\mathbf{A} + 2\mathbf{B}$, (ii) $2\mathbf{B} - 3\mathbf{C}$, (iii) $2\mathbf{A} - \frac{1}{2}(\mathbf{B} + \mathbf{C})$

From the following matrix equations, determine the values of x and y.

$$5. \quad \begin{pmatrix} x+2y & 14 \\ -3 & y-2 \end{pmatrix} = \begin{pmatrix} 4 & 14 \\ -3 & 7+3x \end{pmatrix}$$

$$6. \quad \begin{pmatrix} 1 & x^2+12 \\ x+4 & 3 \end{pmatrix} = \begin{pmatrix} 1 & 7x \\ x+4 & 3 \end{pmatrix}$$

$$7. \quad \begin{pmatrix} 4x^2 & 5 \\ 3 & y^2-4 \end{pmatrix} = \begin{pmatrix} 16 & 5 \\ 3 & 0 \end{pmatrix}$$

In the following matrix equations, find the value of x.

$$8. \quad \begin{pmatrix} 2x & 3 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} -x & 1 \\ 1 & 0 \end{pmatrix} + \begin{pmatrix} 6 & 2 \\ 0 & 0 \end{pmatrix}$$

$$9. \quad \begin{pmatrix} x^2 & 1 \\ 1 & 3 \end{pmatrix} + \begin{pmatrix} 2x & 3 \\ 2 & 4 \end{pmatrix} = \begin{pmatrix} 3 & 4 \\ 3 & 7 \end{pmatrix}$$

10. Find the values of a, b, c and d if:

$$\begin{pmatrix} 2 & 5 \\ 3 & a \end{pmatrix} - \begin{pmatrix} b & c \\ 3 & 4 \end{pmatrix} = \begin{pmatrix} 1 & 9 \\ d & 4 \end{pmatrix}$$

11. If $\mathbf{A} = \begin{pmatrix} x & 3 & -7 \\ 4 & 0 & 6y \end{pmatrix}$ and $\mathbf{B} = \begin{pmatrix} 1-y & 9 & -21 \\ 12 & 0 & 36 \end{pmatrix}$ and $3\mathbf{A} = \mathbf{B}$, find the values of x and y .
12. If $\mathbf{A} = \begin{pmatrix} 2 & 3 \\ 1 & 4 \end{pmatrix}$ and $\mathbf{B} = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}$, find:
 (a) \mathbf{A}^2 , (b) $2\mathbf{AB}$
 (c) $\mathbf{A}(\mathbf{A} + \mathbf{B})$.
13. If $\begin{pmatrix} x & 4 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} -2 \\ 3 \end{pmatrix} = \begin{pmatrix} 4 \\ 0 \end{pmatrix}$, find the value of x .
14. If $\mathbf{K} = \begin{pmatrix} 2 & 3 \\ 5 & 8 \end{pmatrix}$ and $\mathbf{L} = \begin{pmatrix} 8 & -3 \\ -5 & 2 \end{pmatrix}$,
 (a) find \mathbf{IL} and \mathbf{LI} ,
 (b) find $\mathbf{I}^2\mathbf{K}$ and \mathbf{KI}^2 .
15. (a) $\begin{pmatrix} 2 & 3 & 5 \end{pmatrix} \mathbf{K} = \begin{pmatrix} -1 & 0 & 8 & 2 & 4 \end{pmatrix}$. What is the order of matrix \mathbf{K} ?
 (b) If \mathbf{A} is a 3×4 matrix and \mathbf{C} is a matrix such that $\mathbf{A} + \mathbf{B} = \mathbf{C}$, what is the order of \mathbf{B} and of \mathbf{C} ?
16. (a) If a 2×5 matrix is multiplied by a 5×3 matrix, what is the order of the product?
 (b) If \mathbf{A} is a 3×5 matrix and \mathbf{B} is a matrix such that \mathbf{AB} and \mathbf{BA} are both possible, what is the order of \mathbf{B} ?
17. If $\mathbf{A} = \begin{pmatrix} 1 & 1 \\ x & y \end{pmatrix}$ and $\mathbf{A}^2 = \mathbf{I}$, find the values of x and y .
18. If the matrices $\mathbf{P} = \begin{pmatrix} 1 & 3 \\ 4 & 2 \end{pmatrix}$ and $\mathbf{Q} = \begin{pmatrix} 5 & 1 \\ -1 & 0 \end{pmatrix}$
 (a) $\mathbf{PQ} - \mathbf{QP}$, (b) $(\mathbf{P} + \mathbf{Q})^2$
19. If $\begin{pmatrix} x & -6 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ 1 \end{pmatrix} = \begin{pmatrix} x \\ y \end{pmatrix}$, find the possible values of x and y .
20. In an athletics meet in a school, the Blue House got 3 firsts, 4 seconds and nil third in track events and 1 first, 2 seconds and 1 third in relays. The point system is as follows:

	First	Second	Third
Track	3	2	1
Relay	7	5	2

Give this information in the form of two matrices. Multiply the matrices together and label rows and columns of your answer to indicate what the entries represent.

21. Three ladies P, Q and R went for shopping. P bought 2 kg of sugar, $\frac{1}{4}$ kg of tea and 1 loaf of bread; Q bought 10 kg of sugar, $\frac{1}{2}$ kg of tea and 2 loaves of bread; R bought 5 kg of sugar, $\frac{1}{4}$ kg of tea and 1 loaf of bread.

Write this information in the form of a 3×3 matrix.

The prices are: Sugar sh. 1,500 per kg, tea sh. 2,000 per kg and sh. 1,000 for one loaf of bread. Use matrix multiplication to find the total amount of money spent by each of P, Q and R. Label rows and columns of your answer to indicate what the entries represent.

22. The results of four soccer teams are shown below together with a matrix showing how points are awarded.

$$\begin{array}{c}
 \mathbf{W} \quad \mathbf{D} \quad \mathbf{L} \\
 \mathbf{A} \quad \begin{pmatrix} 7 & 3 & 5 \end{pmatrix} \\
 \mathbf{B} \quad \begin{pmatrix} 5 & 4 & 2 \end{pmatrix} \\
 \mathbf{C} \quad \begin{pmatrix} 8 & 2 & 5 \end{pmatrix} \\
 \mathbf{D} \quad \begin{pmatrix} 6 & 3 & 4 \end{pmatrix}
 \end{array}$$

$$\begin{array}{c}
 \text{Points} \\
 \mathbf{W} \quad \begin{pmatrix} 4 \end{pmatrix} \\
 \mathbf{D} \quad \begin{pmatrix} 1 \end{pmatrix} \\
 \mathbf{L} \quad \begin{pmatrix} 0 \end{pmatrix}
 \end{array}$$

Multiply the matrices and hence state which team has most points.

23. A restaurant sells three meals A, B and C. In two days the sales were as shown in matrix **S**.

$$\mathbf{S} = \begin{array}{c} \\ \mathbf{A} \\ \mathbf{B} \\ \mathbf{C} \end{array} \begin{array}{cc} \text{Mon} & \text{Tue} \\ \begin{pmatrix} 10 & 5 \\ 15 & 20 \\ 10 & 10 \end{pmatrix} \end{array}$$

The price in shillings paid for each meal is given by matrix **P**.

$$P = \begin{pmatrix} A & B & C \\ 2000 & 4000 & 7000 \end{pmatrix}$$

Work out the matrix product **PS** and interpret the result.

The Determinant of a 2×2 Matrix

The determinant of $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ is the scalar $ad - bc$.

It is denoted by $\det M$ or $\begin{vmatrix} a & b \\ c & d \end{vmatrix}$.

Note: The elements a and d are in the major or leading diagonal while b and c are in the minor diagonal.

Determinant of a 2×2 matrix = Product of elements in major diagonal *minus* the product of elements in the minor diagonal.

The inverse of a 2×2 matrix

The inverse of a matrix $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, written as M^{-1} , is given by

$$M^{-1} = \frac{1}{\det M} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$$

The matrix $\begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$ is called the adjoint of matrix M .

The adjoint of M is obtained by interchanging the elements in the major diagonal and changing the signs of the elements in the minor diagonal.

Note:

- (i) Given a matrix M , if $\det M = 0$ then M^{-1} cannot be found. In this case M is said to be **singular**.
- (ii) If M^{-1} is the inverse of M , then $MM^{-1} = M^{-1}M = I$.

Example 2.7

Find the inverse of $A = \begin{pmatrix} 1 & 2 \\ 4 & 11 \end{pmatrix}$.

Solution

Change 11 and 1 over and change the signs of 2 and 4 to obtain,

$$\begin{pmatrix} 11 & -2 \\ -4 & 1 \end{pmatrix}$$

Find the determinant of A . That is, $\det A = (11 \times 1) - (4 \times 2) = 11 - 8 = 3$.

$$\text{Then } A^{-1} = \frac{1}{3} \begin{pmatrix} 11 & -2 \\ -4 & 1 \end{pmatrix} = \begin{pmatrix} \frac{11}{3} & -\frac{2}{3} \\ -\frac{4}{3} & \frac{1}{3} \end{pmatrix}.$$

Example 2.8

Calculate the determinant and hence the inverse of the following matrices where possible.

(a) $\begin{pmatrix} 4 & 2 \\ 2 & 1 \end{pmatrix}$

(b) $\begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}$

Solution

(a) $\begin{vmatrix} 4 & 2 \\ 2 & 1 \end{vmatrix} = 4 \times 1 - 2 \times 2$
 $= 4 - 4 = 0$

Since the determinant is 0 the inverse cannot be found.

(b) $\begin{vmatrix} 1 & 1 \\ -1 & 1 \end{vmatrix} = 1 \times 1 - (-1) \times 1$
 $= 1 + 1 = 2$

Inverse of $\begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}$ is $\frac{1}{2} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} \frac{1}{2} & -\frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{pmatrix}$.

Note: If **A**, **B** and **C** are matrices such that **AB = C**, then

$$\mathbf{A}^{-1}\mathbf{AB} = \mathbf{A}^{-1}\mathbf{C}$$

$$\mathbf{IB} = \mathbf{A}^{-1}\mathbf{C}, \text{ since } \mathbf{A}^{-1}\mathbf{A} = \mathbf{I}$$

$$\mathbf{B} = \mathbf{A}^{-1}\mathbf{C} \text{ (} \mathbf{IB} = \mathbf{B} \text{)}.$$

Example 2.9

If $\mathbf{B} = \begin{pmatrix} 3 & -4 \\ 1 & -2 \end{pmatrix}$ and $\mathbf{AB} = \mathbf{I}$, find **A**.

Solution

Since $\mathbf{AB} = \mathbf{I} \Rightarrow \mathbf{A}$ is the inverse of **B**.

$$\begin{vmatrix} 3 & -4 \\ 1 & -2 \end{vmatrix} = 3 \times (-2) - 1 \times (-4)$$
$$= -6 - (-4) = -6 + 4 = -2$$

$$\mathbf{B}^{-1} = -\frac{1}{2} \begin{pmatrix} -2 & 4 \\ -1 & 3 \end{pmatrix} = \begin{pmatrix} 1 & -2 \\ \frac{1}{2} & -\frac{3}{2} \end{pmatrix}$$

Therefore, $\mathbf{A} = \begin{pmatrix} 1 & -2 \\ \frac{1}{2} & -\frac{3}{2} \end{pmatrix}$.

Solution of simultaneous equations by matrix method

A pair of simultaneous equations can also be written in matrix form to obtain the

required solutions.

Example 2.10

Solve the following pair of simultaneous equations:

$$2x - 3y = 7; 4x + 3y = 5$$

Solution

Expressing the equations in matrix form we have,

$$\mathbf{M} \begin{pmatrix} 2 & -3 \\ 4 & 3 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 7 \\ 5 \end{pmatrix} \text{ where } \mathbf{M} \text{ is the matrix of the coefficients.}$$

Write down the inverse of M which is $\frac{1}{18} \begin{pmatrix} 3 & 3 \\ -4 & 2 \end{pmatrix}$

Premultiply both sides of the equation by $\begin{pmatrix} 3 & 3 \\ -4 & 2 \end{pmatrix}$. We need not multiply by $\frac{1}{18}$

as multiplying both sides by $\frac{1}{18}$ does not alter the equation.

$$\begin{pmatrix} 3 & 3 \\ -4 & 2 \end{pmatrix} \begin{pmatrix} 2 & -3 \\ 4 & 3 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 3 & 3 \\ -4 & 2 \end{pmatrix} \begin{pmatrix} 7 \\ 5 \end{pmatrix}$$

$$\begin{pmatrix} 18 & 0 \\ 0 & 18 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 36 \\ -18 \end{pmatrix}$$

$$18 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = 18 \begin{pmatrix} 2 \\ -1 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 2 \\ -1 \end{pmatrix}$$

$$\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 2 \\ -1 \end{pmatrix}$$

$$\text{or } \begin{pmatrix} 18x \\ 18y \end{pmatrix} = \begin{pmatrix} 36 \\ -18 \end{pmatrix}$$

$$18x = 36 \Rightarrow x = 2$$

$$18y = -18 \Rightarrow y = -1$$

$\therefore x = 2$ and $y = -1$

Exercise 2.2

Write down the inverse of the following matrices where possible.

1. $\begin{pmatrix} 2 & 6 \\ 3 & 9 \end{pmatrix}$

2. $\begin{pmatrix} 5 & 7 \\ 2 & 3 \end{pmatrix}$

3. $\begin{pmatrix} -1 & \frac{1}{2} \\ -2 & -1 \end{pmatrix}$

4. $\begin{pmatrix} -2 & 3 \\ -3 & 4 \end{pmatrix}$

5. $\begin{pmatrix} 4 & 0 \\ 0 & \pm 1 \end{pmatrix}$

6. $\begin{pmatrix} 2 & -4 \\ 3 & -6 \end{pmatrix}$

7. Find the inverse of the matrix $\mathbf{A} = \begin{pmatrix} 5 & 1 \\ 2 & 3 \end{pmatrix}$ and show that $\mathbf{AA}^{-1} = \mathbf{A}^{-1}\mathbf{A} = \mathbf{I}$, where \mathbf{I} is the identity matrix.
8. Given $\mathbf{P} = \begin{pmatrix} 3 & -4 \\ -2 & 3 \end{pmatrix}$, find matrix \mathbf{A} , given that $\mathbf{AP} = \mathbf{I}$, where \mathbf{I} is the identity matrix.

Find the values of x and y in Q 9 - 12:

9. $\begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 2 & 2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix}$

10. $\begin{pmatrix} x & y \end{pmatrix} \begin{pmatrix} 2 & 3 \\ 1 & 2 \end{pmatrix} = \begin{pmatrix} 3 & 5 \end{pmatrix}$

11. $\begin{pmatrix} 2 & 3 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 7 \\ 4 \end{pmatrix} = \begin{pmatrix} x \\ y \end{pmatrix}$

12. $\begin{pmatrix} 1 & 2 \\ 2 & 4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$

13. Find the inverse of $\begin{pmatrix} 5 & 3 \\ 2 & 2 \end{pmatrix}$ and hence solve the simultaneous equations:
 $5x + 3y = 7$, $2x + 2y = 2$.

14. $\mathbf{A} = \begin{pmatrix} 5 & 4 \\ 3 & 2 \end{pmatrix}$ and $\mathbf{B} = \begin{pmatrix} 6 & 3 \\ 8 & 4 \end{pmatrix}$.

- (a) Calculate the determinant of the matrices \mathbf{A} and \mathbf{B} .
 (b) Which of the matrices has an inverse? What is the inverse in this case?

15. Given $\mathbf{P} = \begin{pmatrix} a & 2a \\ a-1 & a+1 \end{pmatrix}$ is a singular matrix, find the possible values of a .

16. $\mathbf{P} = \begin{pmatrix} 2 & 1 \\ 3 & 1 \end{pmatrix}$ and $\mathbf{Q} = \begin{pmatrix} -1 & 1 \\ 2 & 0 \end{pmatrix}$. Find the determinant in each case. Find \mathbf{PQ} and hence show that $\det. (\mathbf{PQ}) = \det. (\mathbf{P}) \det. (\mathbf{Q})$.

17. $\mathbf{A} = \begin{pmatrix} 3 & 0 \\ 0 & 4 \end{pmatrix}$, $\mathbf{B} = \begin{pmatrix} a & b \\ 0 & c \end{pmatrix}$ and $\mathbf{AB} = \mathbf{A} + \mathbf{B}$. Find the values of a , b and c .

18. \mathbf{M} is the matrix $\begin{pmatrix} 3 & 2 \\ 4 & 5 \end{pmatrix}$ and \mathbf{P} is the matrix $\begin{pmatrix} 5 \\ 2 \end{pmatrix}$. Calculate the matrices

\mathbf{M}^{-1} and $\mathbf{M}^{-1}\mathbf{P}$. Hence solve the equations:

$$3x + 2y = 5$$

$$4x + 5y = 2.$$

19. Given that $\mathbf{P} = \begin{pmatrix} 2 & 3 \\ 1 & 2 \end{pmatrix}$ and $\mathbf{Q} = \begin{pmatrix} 2 & -3 \\ -1 & 2 \end{pmatrix}$, find the matrix product \mathbf{PQ} and $\mathbf{P}^{-1}\mathbf{Q}$.

A matrix \mathbf{R} of order 2×2 is such that $\mathbf{QR} = \mathbf{P}$. Find the matrix \mathbf{R} .

20. The determinant of matrix \mathbf{A} is 2 and $\mathbf{A}^{-1} = \begin{pmatrix} 1 & -1 \\ -1 & 2 \end{pmatrix}$. Find \mathbf{A} and show

that $\mathbf{AA}^{-1} = \mathbf{I}$.

21. (a) Express the matrix equation $\begin{pmatrix} 5 & 6 \\ 3 & 4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} a \\ b \end{pmatrix}$ into two simultaneous equations.
 (b) Express, as a matrix equation, the two simultaneous equations $5x + 6y = 25$ and $3x + 4y = 17$.
 (c) Hence or otherwise, find the values of x and y .

Solve the following pairs of simultaneous equations using matrix method:

22. $2x + 3y = 7$
 $x + 2y = 3$

23. $5a - 12b = 4$
 $4a - 7b = -2$

24. $3x - 4y = 7$
 $4x + 6y = 15$

25. $2p + 3q = 4$
 $4p + q = -2$

26. $-r + t = 1$
 $3r - 4t = -3$

27. $4x + y = 0$
 $6x - y = 5$

3. Matrices and Transformations

In Book 2, we discussed four types of transformations. These were translations, reflections, enlargements and rotations. There are two other transformations, namely **shears** and **stretches**. Each transformation has different properties on a Cartesian plane.

In this topic we shall consider transformations of the Cartesian plane and the connection between matrices and transformations.

Translations

A translation is a transformation such that the points in the object can be joined to the corresponding points in the image by a set of straight lines which are all equal in length and parallel.

A translation is described by the direction and magnitude of movement (vector). Under a translation, all the points in the object move through the same distance and in the same direction. In order to obtain the image of a point under a translation we add the vector of translation to the position vector of the point.

Example 3.1

Triangle ABC is such that A(-4, 3), B(-1, 1) and C(-2, 5). Find the coordinates of its image after a translation whose vector is $\begin{pmatrix} 6 \\ 2 \end{pmatrix}$

Solution


$$OA_1 = \begin{pmatrix} -4 \\ 3 \end{pmatrix} + \begin{pmatrix} 6 \\ 2 \end{pmatrix} = \begin{pmatrix} 2 \\ 5 \end{pmatrix}$$

$$OB_1 = \begin{pmatrix} -1 \\ 1 \end{pmatrix} + \begin{pmatrix} 6 \\ 2 \end{pmatrix} = \begin{pmatrix} 5 \\ 3 \end{pmatrix}$$

$$OC_1 = \begin{pmatrix} -2 \\ 5 \end{pmatrix} + \begin{pmatrix} 6 \\ 2 \end{pmatrix} = \begin{pmatrix} 4 \\ 7 \end{pmatrix}$$

Therefore the coordinates of the images are $A_1(2, 5)$, $B_1(5, 3)$ and $C_1(4, 7)$.

This can be illustrated on the Cartesian plane as shown in the following diagram.



Rotations

A rotation is a transformation of an object about a fixed point such that every point in the object turns through the same angle relative to the fixed point.

A rotation is described by the angle and the centre of rotation. An anticlockwise rotation is positive while a clockwise rotation is negative.

When an object undergoes a rotation:

- a point and its image are equidistant from the centre of rotation.
- each point of an object moves along an arc of a circle whose centre is the centre of rotation.
- only the centre of rotation remains fixed.
- the perpendicular bisector of a line joining a point and its image passes through the centre of rotation.
- the object and its image are directly congruent.


Example 3.2

A triangle has A(2, 2), B(2, 5) and C(4, 5) as its vertices. Find the coordinates of its image when rotated through 90° anticlockwise about the origin.

Solution

In order to find the coordinates of the image, join O to A to make line OA. From

line OA measure an angle 90° and mark point A_1 such that $OA = OA_1$. Repeat this process with points B and C to get points B_1 and C_1 . Join points A_1 , B_1 and C_1 to get the image of triangle ABC. The coordinates of the vertices of the image are $A_1(-2, 2)$, $B_1(-5, 2)$ and $C_1(-5, 4)$.



Reflections

Under a reflection:

- a point and its image are equidistant from the mirror line.
- the mirror line is a perpendicular bisector of the line joining a point and its image.
- points on the mirror line are invariant.
- an object and its image are indirectly congruent.

In the Cartesian plane, a reflection is described by stating the *equation of the mirror line*.


Example 3.3

Triangle PQR is such that $P(-2, 1)$, $Q(-3, 3)$ and $R(2, 2)$ are the vertices. Find the coordinates of its image under a reflection in the line $x = 2$.

Solution

When the triangle is reflected on the line $x = 2$, the coordinates of the image are $P_1(6, 1)$, $Q_1(7, 5)$ and $R_1(2, 2)$.

R lies on the mirror line and is invariant



Enlargements

Enlargement is described by stating the *centre of enlargement* and the *scale factor*. The enlargement scale factor can be negative or positive.

The ratio $\frac{\text{image distance from centre}}{\text{object distance from centre}}$ is called the **scale factor**.

Example 3.4


Triangle ABC is such that A(2, 1), B(4, 1) and C(3, 4) are its vertices. Find the coordinates of its image after an enlargement with centre (0, 0) and scale factor:

- (a) 2 (b) -2.

Solution

- (a) When the triangle is enlarged about (0, 0) with scale factor of 2, the coordinates of the images are $A_1(4, 2)$, $B_1(8, 2)$ and $C_1(6, 8)$.
- (b) When the triangle is enlarged about (0, 0) with scale factor of -2, the coordinates of the image are $A_2(-4, -2)$, $B_2(-8, -2)$ and $C_2(-6, -8)$.

Under an enlargement, an object and its image are similar but not congruent..



Shears


A shear is a transformation that keeps one line invariant and moves all the other points parallel to this invariant line. The distance moved by each point is proportional to its distance from the invariant line. The constant of this proportionality is called the **shear factor**. The area of the object remains constant.

The shear factor is given by the ratio:

$$\frac{\text{distance moved by a point}}{\text{perpendicular distance of the point from the invariant line}}$$

Example 3.5

In the figure below, rectangle OABC is mapped onto parallelogram OAB₁C₁ under a shear.



Stretches


Properties of a stretch

- The direction of the stretch is perpendicular to the invariant line.
- The scale factor is given by

$$\frac{\text{distance of the image point from the invariant line}}{\text{distance of the object point from the invariant line}}$$
- Points on the opposite sides of the invariant line move in opposite directions but perpendicular to the invariant line.

Example 3.6

The square OABC is stretched, parallel to the X-axis, to become the rectangle OA'B'C'. Such that $OA' = 4 \times OA$



Transformation matrices

Matrices can be used to represent transformations. In the x-y plane, we use 2×2 matrices to represent transformations. The matrices can be determined by the use of the identity matrix or calculations.

The identity matrix

Consider the unit square OIKJ with coordinates O(0, 0), I(1, 0), K(1, 1) and J(0, 1). When the position vectors $I = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $J = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ are written in matrix form,

they give the identity matrix $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. In this matrix, the first column is the position vector of I and the second column is the position vector of J.

Consider a reflection of the unit square in the x-axis. Point I(1, 0) is mapped onto $I_1(1, 0)$. Point J(0, 1) is mapped onto $J_1(0, -1)$. Point O(0, 0) is mapped onto $O_1(0, 0)$ and point K(1, 1) is mapped onto $K_1(-1, -1)$.

The position vectors of $I_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $J_1 = \begin{pmatrix} 0 \\ -1 \end{pmatrix}$ form the matrix $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$

When the position vectors of the unit square are premultiplied by $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ we get,

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix} = \begin{pmatrix} O_1 & I_1 & J_1 & K_1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & -1 & -1 \end{pmatrix}$$


The result is the matrix formed by the position vectors of the images of the unit square under reflection in the x-axis. This means that $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ represents a reflection in the x-axis.

In order to determine a matrix of transformation, we need to know the kind of transformation it is, find the images of points I and J under that transformation, and then write their position vectors in matrix form.

Example 3.7

A triangle with vertices P(2, 2), Q(5, 1) and R(5, 6) is rotated through 90° anti-clockwise about the origin. Find the transformation matrix and the coordinates of its image.

Solution



The images of I and J under this transformation are $I^1(0,1)$ and $J^1(-1, 0)$.

Therefore, the matrix of transformation is $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$.

When we premultiply the matrix resulting from the position vectors of points P, Q and R by the matrix of transformation, we get:

$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 2 & 5 & 5 \\ 2 & 1 & 6 \end{pmatrix} = \begin{pmatrix} P^1 & Q^1 & R^1 \\ 2 & 1 & 6 \\ 2 & 5 & 5 \end{pmatrix}$$

Therefore, the coordinates of the image are $P^1(-2, 2)$, $Q^1(-1, 5)$ and $R^1(-6, 5)$.

We can find the image of any point in the plane by pre-multiplying the position vector of that point by the matrix of the transformation.

Example 3.8

Find the images of the points with the following position vectors under the

transformation whose matrix is $\begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$

(a) $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$

(b) $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$

(c) $\begin{pmatrix} 2 \\ 5 \end{pmatrix}$

(d) $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$

Solution

(a) $\begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 2 \times 1 + 1 \times 0 \\ 1 \times 1 + 2 \times 0 \end{pmatrix} = \begin{pmatrix} 2+0 \\ 1+0 \end{pmatrix} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$

So the image of $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ is $\begin{pmatrix} 2 \\ 1 \end{pmatrix}$

$$(b) \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} = \begin{pmatrix} 0+1 \\ 0+2 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}.$$

So the image of $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ is $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$.

$$(c) \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 2 \\ 5 \end{pmatrix} = \begin{pmatrix} 4+5 \\ 2+10 \end{pmatrix} = \begin{pmatrix} 9 \\ 12 \end{pmatrix}$$

So the image of $\begin{pmatrix} 2 \\ 5 \end{pmatrix}$ is $\begin{pmatrix} 9 \\ 12 \end{pmatrix}$.

$$(d) \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 2+1 \\ 1+2 \end{pmatrix} = \begin{pmatrix} 3 \\ 3 \end{pmatrix}. \text{ So the image of } \begin{pmatrix} 1 \\ 1 \end{pmatrix} \text{ is } \begin{pmatrix} 3 \\ 3 \end{pmatrix}.$$

Parts (a) and (b) above are especially interesting as the images of $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ are the columns of the transformation matrix.

In general,

The matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ transforms the vectors $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ as follows

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix} \rightarrow \begin{pmatrix} a \\ c \end{pmatrix} \text{ and } \begin{pmatrix} 0 \\ 1 \end{pmatrix} \rightarrow \begin{pmatrix} b \\ d \end{pmatrix}$$

Example 3.9


A is a reflection in the line $y = x$. **B** is a reflection in the y -axis. Find the matrix which represents:

(a) **A**

(b) **B**

Solution

(a)




The images of $I(1, 0)$ and $J(0, 1)$ under this transformation (**A**) are $I'(0, 1)$ and



$J^1(1, 0)$.

Therefore, the matrix representing the transformation is $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$

(b)




The images of I and J under transformation **B** are $I'(-1, 0)$ and $J'(0, 1)$,

respectively. Therefore, the matrix representing **B** is $\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$

Example 3.10

Find the matrix which represents an enlargement of scale factor 2 with centre O

Solution



The images of I and J are $I'(2, 0)$ and $J'(0, 2)$, respectively. Therefore, the matrix representing an enlargement of scale factor 2 about O is

$$\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$$

In general, an enlargement of scale factor k with O as centre of enlargement is represented by the matrix $\begin{pmatrix} k & 0 \\ 0 & k \end{pmatrix}$

Note: Finding the matrix of transformation using the above method (based on the images of points I and J) works only when the transformation is a reflection, rotation, enlargement, shear or stretch in which the origin remains fixed.

Finding the matrix of transformation by calculation.

Example 3.11

Triangle ABC with vertices $A(2, 1)$, $B(2, 3)$ and $C(3, 1)$ maps onto $A_1(4, 1)$, $B_1(8, 3)$ and $C_1(5, 1)$ under a shear with the x -axis as the invariant line and shear factor 2. Find the matrix representing this transformation.

Solution

Let the matrix of transformation be $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$.

$$\text{Therefore, } \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} A & B & C \\ 2 & 2 & 3 \\ 1 & 3 & 1 \end{pmatrix} = \begin{pmatrix} A_1 & B_1 & C_1 \\ 4 & 8 & 5 \\ 1 & 3 & 1 \end{pmatrix}$$

Multiplying these matrices gives,

$$\begin{pmatrix} 2a+b & 2a+3b & 3a+b \\ 2c+d & 2c+3d & 3c+d \end{pmatrix} = \begin{pmatrix} 4 & 8 & 5 \\ 1 & 3 & 1 \end{pmatrix}$$

Equating the corresponding elements in the two matrices gives:

$$\begin{array}{ll} 2a + b = 4 & \text{.....(i)} \\ 2a + 3b = 8 & \text{.....(ii)} \\ 3a + b = 5 & \text{.....(iii)} \end{array} \qquad \begin{array}{ll} 2c + d = 1 & \text{.....(iv)} \\ 2c + 3d = 3 & \text{.....(v)} \\ 3c + d = 1 & \text{... ..(vi)} \end{array}$$

Solving equations (i) and (ii) simultaneously gives, $b = 2$ and $a = 1$.

Solving equations (iv) and (v) simultaneously gives, $d = 1$ and $c = 0$.

$$\text{Therefore, } \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$$

The matrix $\begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$ represents a shear of factor 2 with the x -axis invariant.

Example 3.12

A line PQ , in which $P(10, 4)$ and $Q(2, 8)$, is mapped onto the line P_1Q_1 , such that $P_1(5, 2)$ and $Q_1(1, 4)$ after an enlargement of scale factor $\frac{1}{2}$ with centre O . Determine the matrix representing this transformation.

Solution

Let the matrix of transformation be $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 10 & 2 \\ 4 & 8 \end{pmatrix} = \begin{pmatrix} 5 & 1 \\ 2 & 4 \end{pmatrix}$$

Multiplying these matrices and equating the corresponding elements gives,

$$10a + 4b = 5 \dots(i)$$

$$2a + 8b = 1 \dots(ii)$$

$$10c + 4d = 2 \dots(iii)$$

$$2c + 8d = 4 \dots(iv)$$

Solving equations (i) and (ii) simultaneously gives, $a = \frac{1}{2}$, $b = 0$,

Solving equations (iii) and (iv) simultaneously gives $c = \frac{1}{2}$ and $d = 0$.

The matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{pmatrix}$

Therefore $\begin{pmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{pmatrix}$ represents an enlargement of scale factor $\frac{1}{2}$ about the origin.

Describing a matrix using the points I(1, 0) and J(0, 1).

It is possible to describe a transformation in matrix form by considering the effect on the points I(1, 0) and J(0, 1).

We let $I = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $J = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$.

The *columns* of a matrix give us the images of I and J after the transformation.


Example 3.13

Describe the transformation with matrix $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$.

Column $\begin{pmatrix} 0 \\ -1 \end{pmatrix}$ represents I^1 (the image of I).

Column $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ represents J^1 (the image of J)

Draw I, J, I^1 and J^1 on a diagram. Clearly both I and J have been rotated 90° clockwise about the origin.



$\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ represents a rotation of -90° about the origin.

This method can be used to describe a reflection, rotation, enlargement, shear or stretch in which the origin remains fixed.

Exercise 3.1

- Triangle XYZ has vertices X(1, 1), Y(2, 4) and Z(4, 4). Determine the matrix which represents each of the following transformations and hence the coordinates of the vertices of the image triangle $X_1Y_1Z_1$.
 - Reflection in the x-axis
 - Reflection in the y-axis.
 - A rotation of -90° about the origin.
 - A rotation of 90° about the origin.
 - A half turn about the origin.
 - An enlargement of scale factor -2 with centre O.
- Determine the matrix that can transform each of the following pairs of points onto their respective images.
 - A(1, 2) and B(3, 0) onto $A_1(4, 4)$ and $B_1(3, 9)$.
 - A(-2, 1) and B(5, -3) onto $A_1(1, -2)$ and $B_1(1, -3, 5)$
 - M(4, 5) and N(-1, 6) onto $M_1(-4, 5)$ and $N_1(1, 6)$
 - P(8, 3) and Q(0, 5) onto $P_1(8, 3)$ and $Q_1(0, 5)$
- Determine the matrix that can transform each of the following shapes onto their corresponding images.
 - Triangle ABC with vertices A(2, 3), B(2, 6) and C(4, 3) onto triangle $A_1B_1C_1$ with vertices at $A_1(6, 6)$, $B_1(6, 12)$ and $C_1(12, 6)$.
 - Rectangle ABCD with vertices A(-3, -1), B(-3, -4), C(-1, -4), D(-1, -1) onto rectangle $A_1B_1C_1D_1$ with vertices $A_1(-3, 1)$,

$B_1(-3, 4)$, $C_1(-1, 4)$ and $D_1(-1, 1)$.

- (c) Triangle PQR with vertices $P(1, 1)$, $Q(2, 4)$ and $R(4, 1)$ onto triangle $P_1Q_1R_1$ with vertices $P_1(0, 0)$, $Q_1(-4, 2)$ and $R_1(4, -2)$.

4. Use the points I and J to describe the transformation represented by each Matrix.

(a) $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$

(b) $\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$

(c) $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$

(d) $\begin{pmatrix} -2 & 0 \\ 0 & -2 \end{pmatrix}$

5. Draw the triangle $A(2, 2)$, $B(6, 2)$, $C(6, 4)$. Find its image under the transformation represented by the following matrices:

(a) $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$

(b) $\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$

(c) $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$

(d) $\begin{pmatrix} \frac{1}{2} & 0 \\ 0 & \frac{1}{2} \end{pmatrix}$

6. Plot the object and image for the following:

(a) Object: $P(4, 2)$, $Q(4, 4)$, $R(0, 4)$; matrix: $\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$

(b) Object: $A(-6, 8)$, $B(-2, 8)$, $C(-2, 6)$; matrix: $\begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$

Describe each as a single transformation.

7. Find and draw the image of the square $(0, 0)$, $(1, 1)$, $(0, 2)$, $(-1, 1)$ under the transformation represented by the matrix

$$\begin{pmatrix} 4 & 3 \\ -3 & -2 \end{pmatrix}.$$

Show that the transformation is a shear and find the equation of the invariant line.

8. Find and draw the image of the unit square $O(0, 0)$, $I(1, 0)$, $K(1, 1)$, $J(0, 1)$ under the transformation represented by the matrix

$$\begin{pmatrix} 3 & 0 \\ 0 & 2 \end{pmatrix}.$$
 This transformation is called a two-way stretch.

Successive transformations

An object can undergo several transformations, one after the other. This is



done such that the image of the preceding transformation becomes the object of the next transformation.

When an object, A , undergoes a transformation R followed by another transformation N , this is shown as $N[R(A)]$ or $NR(A)$.

The first transformation is always indicated to the right of the second transformation.

$RR(A)$ means 'perform transformation R on A and then perform R on the image'. It may be written $R^2(A)$.


Example 3.14

A is a triangle with vertices $(2, 1)$, $(2, 4)$ and $(4, 1)$. R is a rotation of 90° clockwise about the origin and N is a reflection in the y -axis. Find the vertices of the image of A if it undergoes the following transformations:

- (a) $NR(A)$ (b) $RN(A)$

Solution

- (a) The figure below shows triangle A under these transformations and its image $R(A)$ and $NR(A)$.




The final image of A has vertices $(-1, -2)$, $(-1, -4)$ and $(-4, -2)$.

- (b) The figure below shows triangle A and its images $N(A)$ and $RN(A)$.

The final image of triangle A under these transformations has vertices $(1, 2)$, $(1, 4)$ and $(4, 2)$.


The order of transformation is very important.

$NR(A) \neq RN(A)$.



Example 3.15

Triangle B has vertices at $(-2, 1)$, $(-4, 1)$ and $(-4, 4)$. T is a translation represented by the vector $\begin{pmatrix} 8 \\ 3 \end{pmatrix}$, R is a reflection in the line $y = -x$ and M is a rotation of 90° about the origin. Find the vertices of $MRT(B)$.



Exercise 3.2

Draw x - and y -axes with values from -8 to $+8$ and plot the point $P(3, 2)$.

R denotes 90° clockwise rotation about $(0, 0)$;

X denotes reflection in $x = 0$.

H denotes 180° rotation about $(0, 0)$;

T denotes translation $\begin{pmatrix} 3 \\ 2 \end{pmatrix}$.

For each question, write down the coordinates of the final image of P.

- | | |
|-----------------------|-------------------------|
| 1. R(P) | 2. TR(P) |
| 3. T(P) | 4. RT(P) |
| 5. TH(P) | 6. XT(P) |
| 7. HX(P) | 8. XX(P) |
| 9. R ² (P) | 10. T ³ X(P) |

Exercise 3.3

In this exercise, transformations A, B,H, are as follows:

A denotes reflection in $x = 2$

B denotes 180° rotation, centre (1, 1)

C denotes translation $\begin{pmatrix} -6 \\ 2 \end{pmatrix}$

D denotes reflection in the line $y = x$.

E denotes reflection in $y = 0$

F denotes translation $\begin{pmatrix} 4 \\ 3 \end{pmatrix}$

G denotes 90° rotation clockwise, centre (0, 0)

H denotes enlargement, scale factor $\frac{1}{2}$, centre (0, 0).

Draw the axes with values from -8 to +8.

- Draw triangle LMN at L(2, 2), M(6, 2), N(6, 4). Find the image of LMN under the following combinations of transformations. Write down the coordinates of the image of point L in each case:
 - CA(LMN)
 - ED(LMN)
 - DB(LMN)
 - BE(LMN)
 - EB(LMN)
- Draw triangle PQR at P(2, 2), Q(6, 2), R(6, 4). Find the image of PQR under the following combinations of transformations. Write down the coordinates of the image of point P in each case.
 - AF(PQR)
 - CG(PQR)
 - AG(PQR)
 - HE(PQR)
- Draw triangle XYZ at X(-2, 4), Y(-2, 1), Z(-4, 1). Find the image of XYZ under the following combinations of transformations and in each case state

the equivalent single transformation.

- (a) $G^2E(XYZ)$ (b) $CB(XYZ)$
(c) $DA(XYZ)$

4. Draw triangle OPQ at O(0, 0), P(0, 2), Q(3, 2). Find the image of OPQ under the following combinations of transformations and state the equivalent single transformation in each case:

- (a) $DE(OPQ)$ (b) $FC(OPQ)$
(c) $DEC(OPQ)$ (d) $DFE(OPQ)$

5. Draw triangle XYZ at X(1, 2), Y(1, 6), Z(3, 6).

- (a) Find the image of XYZ under each of the transformations **BC** and **CB**.
(b) Describe fully the single transformation equivalent to **BC**.
(c) Describe fully the transformation **M** such that $MCB = BC$.

Matrices of successive transformations

If each of the successive transformations has a transformation matrix, then a single matrix that represents all the transformations can be obtained. Such a matrix maps the original object onto the final image.

Example 3.16

M and **N** are transformations represented by matrices $\mathbf{M} = \begin{pmatrix} 1 & 2 \\ 1 & 3 \end{pmatrix}$ and

$\mathbf{N} = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}$. P is a point with coordinates (-1, 2).

- (a) Find:
(i) $\mathbf{M}(P)$ (ii) $\mathbf{N}(P)$
(iii) $\mathbf{MN}(P)$.
(b) If **R** is a transformation such that $\mathbf{R} = \mathbf{MN}$, find $\mathbf{R}(P)$. What can you deduce from the results of (a) and (b)?

Solutions

(a) (i) $\mathbf{M}(P)$ is $\begin{pmatrix} 1 & 2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} -1 \\ 2 \end{pmatrix} = \begin{pmatrix} 3 \\ 5 \end{pmatrix}$.

$P_1(3, 5)$

(ii) $\mathbf{N}(P)$ is $\begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} -1 \\ 2 \end{pmatrix} = \begin{pmatrix} -1 \\ 0 \end{pmatrix}$

$P_2(-1, 0)$

$$\begin{aligned} \text{(iii)} \quad \mathbf{MN(P)} &= \mathbf{M[N(P)]} = \mathbf{M} \begin{pmatrix} -1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 & 2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} -1 \\ 0 \end{pmatrix} \\ &= \begin{pmatrix} 1 \\ -1 \end{pmatrix} \end{aligned}$$

$$\text{(b)} \quad \mathbf{R} = \mathbf{MN} = \begin{pmatrix} 1 & 2 \\ 1 & 3 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix} = \begin{pmatrix} 5 & 2 \\ 7 & 3 \end{pmatrix}$$

$$\mathbf{R(P)} = \begin{pmatrix} 5 & 2 \\ 7 & 3 \end{pmatrix} \begin{pmatrix} -1 \\ 2 \end{pmatrix} = \begin{pmatrix} -1 \\ -1 \end{pmatrix}$$

From the results of (a) and (b), $\mathbf{MN(P)} = \mathbf{R(P)}$.

Example 3.17

Draw ΔPQR with vertices $P(3, 1)$, $Q(6, 1)$ and $R(6, 5)$.

- (a) Draw the image of ΔPQR under the transformation represented by the matrix $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, label it $P_1Q_1R_1$ and describe this transformation fully.
- (b) Draw the image of $\Delta P_1Q_1R_1$ under a rotation of 180° about the origin and label it $P_2Q_2R_2$. Find the matrix that represents this transformation.
- (c) Describe fully the single transformation that maps ΔPQR onto $\Delta P_2Q_2R_2$. Find the matrix that represents this transformation.

Solutions

$$\text{(a)} \quad \begin{matrix} & P & Q & R & & P_1 & Q_1 & R_1 \\ \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} & \begin{pmatrix} 3 & 6 & 6 \\ 1 & 1 & 5 \end{pmatrix} & = & \begin{pmatrix} 1 & 1 & 5 \\ 3 & 6 & 6 \end{pmatrix} \end{matrix}$$

The transformation is a reflection in the line $y = x$.


- (b) The image of $\Delta P_1Q_1R_1$ is $\Delta P_2Q_2R_2$ as shown in the figure below.
- $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$ is the matrix representing this transformation. This is a half-turn about $(0, 0)$.

- (c) From the diagram, the single transformation that maps ΔPQR onto $\Delta P_2Q_2R_2$ is a reflection in $y = -x$.

Under a reflection in $y = -x$, the identity matrix

$$\begin{matrix} I & J \\ \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \end{matrix} \text{ maps onto } \begin{matrix} I_1 & J_1 \\ \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \end{matrix}. \text{ Therefore, } \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \text{ is the matrix that maps}$$

ΔPQR onto $\Delta P_2Q_2R_2$



Alternatively, this transformation is a successive transformation of a reflection in $y = x$ followed by a rotation of 180° about the origin. Thus, a single matrix is obtained from the product of the two matrices as:

$$\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$$

When multiplying, the matrix of the first transformation (in this case reflection in $y = x$) is to the right of the matrix of the second transformation (in this case rotation of 180° about the origin)

Exercise 3.4

1. Find a single matrix that represents the following successive transformations.
 - (a) A reflection in the x-axis followed by a reflection in the y-axis.
 - (b) A reflection in the line $y = x$ followed by a reflection in the line $y = -x$.
 - (c) A clockwise rotation of 90° about the origin followed by an enlargement of scale factor 3 with the centre of enlargement as the origin.
 - (d) An enlargement of factor 2 with centre O followed by an anticlockwise rotation of 90° about O.

2. Draw the parallelogram formed by the points $P(1, -3)$, $Q(4, -3)$, $R(6, 1)$ and $S(3, -1)$.
- Draw the image $P_1Q_1R_1S_1$ of the parallelogram after a reflection in the line $y = 0$.
 - Reflect the image in the line $y = x$ to obtain parallelogram $P_2Q_2R_2S_2$.
 - Describe fully the single transformation that maps $PQRS$ onto $P_2Q_2R_2S_2$.
 - Find the matrix of transformation that maps $PQRS$ onto $P_2Q_2R_2S_2$.
3. Draw triangle T whose vertices are $(4, 3)$, $(-3, -1)$ and $(-1, 3)$.
- Triangle R is the image of triangle T under the transformation represented by the matrix $\mathbf{N} = \begin{pmatrix} -0.6 & 0.8 \\ 0.8 & 0.6 \end{pmatrix}$
 - Calculate the coordinates of the vertices of triangle R .
 - Draw and label triangle R
 - Describe fully a single transformation that maps triangle T onto triangle R .
4. Triangle XYZ is such that $X(1, 4)$, $Y(1, 7)$ and $Z(3, 1)$ are its vertices. Triangle XYZ is mapped onto triangle $X_1Y_1Z_1$ by the transformation represented by the matrix $\mathbf{M} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.
- Draw and label triangle $X_1Y_1Z_1$ and describe fully the transformation that maps triangle XYZ onto $X_1Y_1Z_1$ in geometrical terms.
 - Reflect $\triangle XYZ$ in the y -axis, and label the vertices of its image as X_2 , Y_2 , and Z_2 . Write down the elements of matrix \mathbf{N} which represents this transformation.
 - Triangle $X_1Y_1Z_1$ can be mapped onto triangle $X_2Y_2Z_2$ by a single transformation \mathbf{P} .
 - Describe this transformation fully.
 - Write down the elements of matrix \mathbf{P} which represents this transformation.
 - State the relationship between \mathbf{M} , \mathbf{N} and \mathbf{P} .

The inverse of a transformation

The inverse of a transformation reverses the transformation, i.e. it is the transformation which takes the image back to the object. For example, the inverse of an anticlockwise rotation of 90° about O is a clockwise rotation of 90° about O .

If translation **T** has vector $\begin{pmatrix} 3 \\ -2 \end{pmatrix}$, the translation which has the opposite effect has vector $\begin{pmatrix} -3 \\ 2 \end{pmatrix}$. This is written as T^{-1} .

If **M** is a transformation representing a transformation, then the matrix representing the inverse of the transformation is indicated as M^{-1} .

Example 3.18

Matrix **N** = $\begin{pmatrix} -1 & -2 \\ 2 & 3 \end{pmatrix}$ describes a transformation on ΔPQR . The coordinates of the image are $P_1(-3, 2)$, $Q_1(2, 1)$ and $R_1(0, 1)$. Find:

- the matrix that maps $\Delta P_1Q_1R_1$ onto ΔPQR .
- the coordinates of P, Q and R.

Solution

- The matrix that maps $\Delta P_1Q_1R_1$ onto ΔPQR is the inverse of **N**.

$$N^{-1} = \frac{1}{1} \begin{pmatrix} 3 & 2 \\ -2 & -1 \end{pmatrix} = \begin{pmatrix} 3 & 2 \\ -2 & -1 \end{pmatrix}$$

$$(b) \begin{matrix} & P_1 & Q_1 & R_1 & & P & Q & R \\ \begin{pmatrix} 3 & 2 \\ -2 & -1 \end{pmatrix} & \begin{pmatrix} -3 & 2 & 0 \\ 2 & 1 & 1 \end{pmatrix} & = & \begin{pmatrix} -5 & 8 & 2 \\ 4 & -5 & -1 \end{pmatrix} \end{matrix}$$

Therefore, the coordinates of PQR are P(-5, 4), Q(8, -5), R(2, -1)

Example 3.19

A(2, 1), B(5, 1) and C(5, 3) are vertices of ΔABC and **M** = $\begin{pmatrix} 2 & 1 \\ 3 & 2 \end{pmatrix}$.

- Find the image $A_1B_1C_1$ of ABC under the transformation represented by matrix **M**.
- Find M^{-1} .
- Find the image of $A_1B_1C_1$ under the transformation represented by matrix M^{-1} .
- Comment on your results in (c).

Solution

$$(a) \begin{matrix} & A & B & C & & A_1 & B_1 & C_1 \\ \begin{pmatrix} 2 & 1 \\ 3 & 2 \end{pmatrix} & \begin{pmatrix} 2 & 5 & 5 \\ 1 & 1 & 3 \end{pmatrix} & = & \begin{pmatrix} 5 & 11 & 13 \\ 8 & 17 & 21 \end{pmatrix} \end{matrix} \quad \text{Therefore, the coordinates of the}$$

vertices of the image are: $A_1(5, 8)$, $B_1(11, 17)$ and $C_1(13, 21)$.

Area and the determinant of a matrix.

Under some transformations, the size and shape of objects do not change, while under other transformations they change. When a matrix represents a transformation, the ratio of the area of the image to the area of the object is equal to the determinant of the matrix of transformation.

Example 3.20


A unit square with vertices $O(0, 0)$, $I(1, 0)$, $K(1, 1)$ and $J(0, 1)$ is given a transformation represented by matrix $\mathbf{N} = \begin{pmatrix} 3 & 0 \\ 0 & 2 \end{pmatrix}$.

- Draw the square and its image.
- Find the determinant of \mathbf{N} .
- Find the area of the image $O_1I_1K_1J_1$.

Solution

$$(a) \begin{matrix} & O & I & K & J & & O & I_1 & K_1 & J_1 \\ \begin{pmatrix} 3 & 0 \\ 0 & 2 \end{pmatrix} & \begin{pmatrix} 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix} & = & \begin{pmatrix} 0 & 3 & 3 & 0 \\ 0 & 0 & 2 & 2 \end{pmatrix} \end{matrix}. \text{ Therefore, the image has vertices at:}$$

$O(0, 0)$, $I_1(3, 0)$, $K_1(3, 2)$ and $J_1(0, 2)$.



- The determinant of \mathbf{N} is $(3 \times 2) - (0 \times 0) = 6$.
- The area of $O_1I_1K_1J_1 = 3 \times 2 = 6$ square units.

The area of the unit square is 1 square unit.

The area of the image is 6 square units. From this information we have

$\frac{\text{the area of the image}}{\text{the area of the object}}$ is the same as the determinant of the transformation

matrix.

- (b) Describe transformation R fully.
6. Points A(-5, 1), B(-1, -1) and C(2, 5) are three vertices of rectangle ABCD.
- Determine the coordinates of D.
 - Describe fully a single transformation (not a reflection) that maps A onto D, and B onto C.
7. Triangle T has vertices (2, 1), (4, 1) and (3, 3).
- Find the area of the triangle.
 - Find the area of the image of T when it is transformed under the matrix:

(i) $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$	(ii) $\begin{pmatrix} 2 & 2 \\ 1 & 3 \end{pmatrix}$
(iii) $\begin{pmatrix} 3 & 4 \\ 1 & 3 \end{pmatrix}$	(iv) $\begin{pmatrix} 6 & 0 \\ 4 & \frac{1}{3} \end{pmatrix}$.
8. Triangle K, whose vertices are P(2, 3), Q(5, 3) and R(4, 1), is mapped onto triangle K_1 whose vertices are $P_1(-4, 3)$, $Q_1(-1, 3)$ and $R_1(x, y)$ by a transformation given by matrix $\mathbf{M} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$.
- Find the:
 - elements of matrix \mathbf{M}
 - coordinates of R_1 .
 - Triangle K_2 is the image of triangle K_1 under a reflection in the line $y = x$. Find a single matrix that maps K onto K_2 .
9. Draw triangle XYZ with X(1, 2), Y(1, 5) and Z(2, 5).
- Draw the image of triangle XYZ under the transformation represented by the matrix $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ and label it $X_1Y_1Z_1$.
 - Describe this transformation.
 - Draw the image of triangle XYZ under a half-turn about the origin and label it $X_2Y_2Z_2$.
 - Find the matrix which represents this transformation.
 - Describe the single transformation that maps triangle $X_1Y_1Z_1$ onto $X_2Y_2Z_2$.
 - Find the matrix which represents this transformation.
10. On a squared paper draw a triangle with its vertices at (3, -1), (5, -1) and (5, -4) and label it L.
- Triangle L is mapped onto triangle L_1 under an anticlockwise rotation of 90° about the origin. Draw L_1 in the same plane.

- (b) Triangle L is mapped onto triangle L_2 under a transformation represented by $\mathbf{M} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. Draw L_2 in the same plane.
- (c) Triangle L_3 has vertices $(-3, -1)$, $(-7, -1)$ and $(-7, 5)$. Draw L_3 in the same plane.
- (d) Describe fully the single transformation that maps triangle L onto L_3 .
11. Draw triangle R whose vertices are $(1, 1)$, $(1, 3)$ and $(5, 3)$. Triangle S is the image of triangle R under the transformation represented by
- $$\mathbf{M} = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}.$$
- (a) Calculate the area of triangle R.
- (b) Describe fully the single transformation that is represented by matrix \mathbf{M} .
- (c) Find \mathbf{M}^{-1} .
- (d) What is the image of triangle R under the transformation represented by \mathbf{M}^{-1} ?
12. Draw and label a triangle whose vertices are $A(1, 4)$, $B(2, 4)$ and $C(2, 1)$.
- (a) Enlargement \mathbf{E} with centre at the origin maps triangle ABC onto triangle $A_1B_1C_1$. Given that the coordinates of A_1 are $(4, 16)$:
- draw and label triangle $A_1B_1C_1$.
 - state the scale factor of \mathbf{E} .
- (b) Point $B_2(-4, -2)$ is the image of B under a reflection in line L. Find the equation of L.
- (c) \mathbf{R} is a clockwise rotation of 90° about the origin and it maps triangle ABC onto triangle $A_3B_3C_3$.
- Draw and label triangle $A_3B_3C_3$.
 - Find the matrix representing \mathbf{R} .
- (d) Transformation \mathbf{S} is represented by the matrix $\begin{pmatrix} -3 & 0 \\ 0 & 1 \end{pmatrix}$ and maps triangle ABC onto triangle $A_4B_4C_4$.
- Find the coordinates of A_4 , B_4 and C_4 .
 - Describe transformation \mathbf{S} fully.
13. Triangle ABC has its vertices at $A(6, 0)$, $B(9, 2)$ and $C(6, 2)$.
- (a) Determine the coordinates of the vertices of its image, $A_1B_1C_1$, after a transformation \mathbf{P} , where \mathbf{P} represents a reflection in the line $y = 3$.
- (b) If \mathbf{Q} represents an anticlockwise quarter-turn about the point $(2, 3)$, determine the coordinates of the vertices of triangle $A_2B_2C_2$, the image of triangle $A_1B_1C_1$ under transformation \mathbf{Q} .

14. The vertices of ΔPQR are $P(1, 1)$, $Q(2, 2)$ and $R(0, 3)$.
- Draw and label ΔPQR .
 - The vertices of $\Delta P_1Q_1R_1$ are found at $P_1(-2, 2)$, $Q_1(-3, 3)$ and $R_1(-4, 1)$. Draw and label $\Delta P_1Q_1R_1$.
 - Describe fully the single transformation that maps ΔPQR onto $\Delta P_1Q_1R_1$.
 - Triangle PQR can also be mapped onto $\Delta P_1Q_1R_1$ by an anticlockwise rotation of 90° about the origin, followed by a translation. Write down the column vector which represents this translation.
15. (a) Draw the axes so that both x and y can take values from -2 to $+8$.
- (b) Draw triangle ABC at $A(2, 1)$, $B(7, 1)$, $C(2, 4)$.
- (c) Find the image of ABC under the transformation represented by the matrix $\begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}$ and plot the image on the graph.
- (d) The transformation is a rotation followed by an enlargement. Calculate the angle of the rotation and the scale factor of the enlargement.
16. (a) On graph paper, draw the triangle T whose vertices are $(2, 2)$, $(6, 2)$ and $(6, 4)$.
- (b) Draw the image U of T under the transformation whose matrix is $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.
- (c) Draw the image V of T under the transformation whose matrix is $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$.
- (e) Describe the single transformation which would map U onto V .
17. (a) Find the images of the points $(1, 0)$, $(2, 1)$, $(3, -1)$, $(-2, 3)$ under the transformation with matrix $\begin{pmatrix} 1 & 3 \\ 2 & 6 \end{pmatrix}$.
- (b) Show that the images lie on a straight line, and find its equation.
18. The transformation with matrix $\begin{pmatrix} 2 & 3 \\ 6 & 9 \end{pmatrix}$ maps every point in the plane onto a line. Find the equation of the line.
19. Using a scale of 1 cm to one unit in each case draw the axes taking values of x from -4 to $+6$ and values of y from 0 to 12 .
- (a) Draw and label the quadrilateral $OABC$ with $O(0, 0)$, $A(2, 0)$, $B(4, 2)$, $C(0, 2)$

- (b) Find and draw the image of OABC under the transformation whose matrix is \mathbf{R} , where $\mathbf{R} = \begin{pmatrix} 2.4 & -1.8 \\ 1.8 & 2.4 \end{pmatrix}$.
- (c) Calculate, in surd form, the lengths OB and O'B'
- (d) Calculate the angle AOA'.
- (e) Given that the transformation \mathbf{R} consists of a rotation about O followed by an enlargement, state the angle of the rotation and the scale factor of the enlargement.

20. The matrix $\mathbf{R} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$ represents a positive rotation of θ° about the origin. Find the matrix which represents a rotation of:
- | | |
|----------------|------------------|
| (a) 90° | (b) 180° |
| (c) 30° | (d) -90° |
| (e) 60° | (f) 150° |
| (g) 45° | (h) 53.1° |

Confirm your results for parts (a), (e), (h) by applying the matrix to the quadrilateral O(0, 0), A(0, 2), B(4, 2), C(4, 0).

21. The image (x', y') of a point (x, y) under a transformation is given by
- $$\begin{pmatrix} x' \\ y' \end{pmatrix} = \begin{pmatrix} 3 & 0 \\ 1 & -2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} + \begin{pmatrix} 2 \\ 5 \end{pmatrix}$$
- (a) Find the coordinates of the image of the point (4, 3).
- (b) The image of the point (m, n) is the point (11, 7). Write down two equations involving m and n and hence find the values of m and n.
- (c) The image of the point (h, k) is the point (5, 10). Find the values of h and k.

22. Draw A(0, 2), B(2, 2), C(0, 4) and its image under an enlargement, A'(2, 2), B'(6, 2), C'(2, 6).
- (a) What is the centre of enlargement?
- (b) Find the image of ABC under a reflection in the line $x = 0$
- (c) Find the translation which maps this image onto A' B' C'.
- (d) What is the matrix \mathbf{X} and vector \mathbf{v} which represents a reflection in the line $x = 2$?

23. $\mathbf{A} = \begin{pmatrix} 2 & 0 \\ 1 & 2 \end{pmatrix}$; h and k are numbers so that $\mathbf{A}^2 = h\mathbf{A} + k\mathbf{I}$, where $\mathbf{I} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$.

Find the values of h and k.

24. $\mathbf{M} = \begin{pmatrix} a & 1 \\ 1 & -a \end{pmatrix}$. Find the values of a if: (a) $\mathbf{M}^2 = 17 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ (b) $|\mathbf{M}| = -10$.

25. (a) Draw the x and y axes from -5 to 5 using a scale of 1 cm to represent 1 unit on each axis.
 Draw triangle ABC with A(1, 1), B(4, 1) and C(4, 2).
- (b) (i) Draw the image of triangle ABC when it is rotated 90° anticlockwise about the origin. Label this image $A_1B_1C_1$.
- (ii) Triangle $A_1B_1C_1$ is translated by the vector $\begin{pmatrix} 3 \\ 1 \end{pmatrix}$. Draw and label this image $A_2B_2C_2$.
- (iii) Describe fully a single transformation which maps triangle ABC onto $A_2B_2C_2$.
- (c) (i) Draw the image of triangle ABC under the transformation represented by the matrix $\begin{pmatrix} 1 & 0 \\ 0 & -2 \end{pmatrix}$. Label this image $A_3B_3C_3$.
- (ii) Describe fully the single transformation which maps triangle ABC onto triangle $A_3B_3C_3$.

4. Statistics

- (a) **Statistics** is that branch of mathematics which is concerned with the collection, organization, interpretation, presentation and analysis of numerical data. To a statistician, any information collected is called **data**.



When data has not been ordered in any specific way after collection, it is called **raw data**.

- (b) **Discrete data**. This is a type of data which can take only exact or integral values. For example, the number of cars passing a check point in 30 minutes; the number of students served in the dining hall in five minutes, etc.
- (c) **Continuous data**. This is data which cannot take exact or integral values, but can be given only within a certain range or measured to a certain degree of accuracy. For example, the heights of students in a school; the time taken by each of a class of students to perform a task, etc.

Methods of presentation of data

Pie Chart.

In a pie chart, each number is represented by the area of a sector of a circle. The frequency is proportional to the angle of the sector.

Example 4.1

The table below shows the number of cars of different colors in a car park. Draw a pie chart to represent this information.

Color	Green	Blue	Yellow	Others
Number	10	14	20	16

The angles of the sectors are calculated as follows:


The total number of cars = $10 + 14 + 20 + 16 = 60$

Angle representing green cars = $\frac{10}{60} \times 360^\circ = 60^\circ$

Angle representing blue cars = $\frac{14}{60} \times 360^\circ = 84^\circ$

Angle representing yellow cars = $\frac{20}{60} \times 360^\circ = 120^\circ$

Angle representing other cars = $\frac{16}{60} \times 360^\circ = 96^\circ$.



Frequency Distributions

(a) Discrete data:

To illustrate data more concisely we count the number of times each value occurs and form a frequency distribution. For example, the following data gives the number of blind students in 10 randomly chosen classes in certain school. 0, 2, 1, 4, 2, 3, 2, 1, 4, 5. this information can be presented as follows:

Number of students	Frequency
0	1
1	2
2	3
3	1
4	2
5	1
Total	10

(b) Continuous data:

In connection with large sets of data, a good overall picture and sufficient information can often be conveyed by grouping the data into a number of classes (intervals).

General rules for grouping data:

- (i) Determine the largest and smallest numbers in the raw data and find the difference between them.
- (ii) Divide this difference into appropriate number of class intervals having the same size - the number of intervals usually taken is between 5 and 20 depending on the size of the data (but preferably 7 to 10 for medium data). If this is not feasible use class intervals of different sizes or open class intervals.
- (iii) Determine the number of observations falling into each class interval i.e. find the class frequencies. This is best done by using tallies.
- (iv) Display the results in the form of a table.

Example 4.2

Construct a distribution of the following data on the length of time (in minutes) it took 80 persons to complete a certain task.

23 24 18 14 20 24 24 26 23 21
 16 15 19 20 22 14 13 20 19 27
 29 22 38 28 34 32 23 19 21 31
 16 28 19 18 12 27 15 21 25 16
 30 17 22 29 29 18 25 20 16 11
 17 12 15 24 25 21 22 17 18 15
 21 20 23 18 17 15 16 26 23 22
 11 16 18 20 23 19 17 15 20 10

Solution

Since the smallest value is 10 and the highest is 38, we might choose the three classes 10-19, 20-29, and 30-39; we might choose the six classes: 10-14, 15-19, 20-24, 25-29, 30-34, and 35-39; to mention a few possibilities. Note that in each case the class intervals accommodate all of the data, they do not overlap, and they are all of the same class size. Taking the second classification, we now tally the 80 observations and get the results shown in the following table.

Minute	Tally	Frequency
10 - 14	### ///	8
15 - 19	### ### ### ### ### ///	28
20 - 24	### ### ### ### ### ///	27
25 - 29	### ### ///	12
30 - 34	////	4
35 - 39	/	1
	total	80

The numbers given in the right hand column of this table, which show how many items fall into each class, are called **class frequencies**. The smallest and largest values that can go into any given class are referred to as its **class limits**, and in our example they are: 10, 14, 15, 19, 20, ..., 34, 35, and 39. More specifically, 10, 15, 20, ... and 35 are called the **lower class limits**, and 14, 19, 24, 29, 34 and 39 are called the **upper class limits**.

The lengths of time which we grouped in our example were all given to the nearest minute, so that the first class actually covers the interval from 9.5 minutes to 14.5 minutes, the second class covers the interval from 14.5 to 19.5, and so forth. These numbers are referred to as **class boundaries** or the "real" class limits.

Class marks are simply the **midpoints** of the classes, and they are obtained by the formula:

$$\text{Class mark} = \frac{\text{lower class lim} + \text{upper class lim}}{2}$$

$$\text{Or } \frac{\text{upper class boundary} + \text{lower class boundary}}{2}$$

A **class interval** (width or size) is the length of a class, or the range of values it can contain and is given by the difference between its class boundaries. If the classes are of equal length, their common class width is also given by the difference between any two successive class marks.

Note that the class widths are **not** given by the differences between the respective class limits.

Cumulative frequency

Cumulative frequency is the total frequency up to a given point.

Example 4.3

Convert the distribution above into a cumulative frequency distribution.

Solution

Since none of the values is less than 10, 8 are less than 15, $8+28 = 36$ are less than 20, $8+28+27 = 63$ are less than 25, ... the results are as shown in the following table.

Minutes	Cumulative freq.
Less than 10	0
Less than 15	8

Less than 20	36
Less than 25	63
Less than 30	75
Less than 35	79
Less than 40	80

Histograms

A histogram is constructed by representing the measurements or observations that are grouped on a horizontal scale, the class frequencies on a vertical scale, and drawing rectangles whose bases equal the class intervals (widths) and whose heights are determined by the corresponding class frequencies. The markings on the horizontal scale are class boundaries.

When class widths are not all equal, the class frequencies are represented by the areas of the rectangles instead of their heights. The vertical axis is not labeled frequency but frequency density.

$$\text{Frequency density} = \frac{\text{frequency}}{\text{class width}}$$

Because the area of the bar represents frequency, the height must be adjusted to correspond with the area of the bar.

Histograms can be used to represent both discrete and continuous data, but their main purpose is for use with continuous data.

Example 4.4

Draw a histogram for the data given below.

Time (Minutes)	Tally	Frequency
10 - 14	### ///	8
15 - 19	### ### ### ### ### ///	28
20 - 24	### ### ### ### ### ///	27
25 - 29	### ### ///	12
30 - 34	///	4
35 - 39	/	1
	total	80


Solution

Time (Minutes)	Class boundaries	Frequency
----------------	------------------	-----------



10 - 14	9.5 - 14.5	8
15 - 19	14.5 - 19.5	28
20 - 24	19.5 - 24.5	27
25 - 29	24.5 - 29.5	12
30 - 34	29.5 - 34.5	4
35 - 39	34.5 - 39.5	1
	total	80

Note that all classes have equal width and therefore, we plot frequency against class boundaries.




Example 4.5

The ages of 120 people who traveled by air on Christmas day were recorded and are shown in the frequency table.

Age (yrs)	Frequency
- 10	18
- 20	46
- 30	35
- 40	13
- 50	8

Solution

The notation '- 10' means ' $0 < \text{age} \leq 10$ ' and similarly '- 20' means ' $10 < \text{age} \leq 20$ '. The class boundaries are 0, 10, 20, 30, 40 and 50. The histogram is then drawn as shown in the diagram below (see next page).



Frequency polygons.


In these, class frequencies (or frequency densities in case of a distribution with unequal class width) are plotted against the class marks and the successive points are connected by means of straight lines. Classes with zero frequencies are added at both ends of the distribution so as to obtain a closed polygon.

Example 4.6


Draw a frequency polygon for the data below.

Weight (kg)	Class marks	Frequency
30 - 40	35	5
40 - 50	45	7
50 - 60	55	10
60 - 70	65	5
70 - 80	75	3

Solution:



A frequency polygon can also be formed by joining the mid-points of the tops of the rectangles in a histogram by straight lines. This is known as 'superimposing' a frequency polygon on a histogram (see diagram below).



Cumulative frequency curve (Ogive)


Cumulative frequency is the total frequency up to a given point. A cumulative frequency curve is obtained by plotting cumulative frequency against the upper class boundaries and the points connected by means of a smooth curve. A class with zero frequency is added at the beginning of the distribution so that the resulting curve starts from the 'origin'.

Example 4.7

The marks obtained by 80 students in an examination are shown below.

Mark	frequency	Cum. frequency	Upper class boundaries
- 0	0	0	0.5
1 - 10	3	3	10.5
11 - 20	5	8	20.5
21 - 30	5	13	30.5
31 - 40	9	22	40.5
41 - 50	11	33	50.5
51 - 60	15	48	60.5
61 - 70	14	62	70.5
71 - 80	8	70	80.5
81 - 90	6	76	90.5
91 - 100	4	80	100.5

The corresponding Ogive is then plotted as shown in the following diagram.



Measures of central tendency

The measures of central tendency are the mean, median and mode. These are Values about which the distribution of a set of data is considered to be roughly balanced.

The mean

The mean is the value obtained when then the total sum of the values of the members in a list or distribution is divided by the total frequency. The mean can be obtained for grouped or ungrouped data.

Example 4.8

On a certain day, nine students received, respectively, 1, 3, 2, 0, 1, 5, 2, 1 and 3 pieces of mail. Find the mean.

Solution

The total number of pieces of mail which these students received is $1+3+2+0+1+5+2+1+3 = 18$, so that the mean number per student is

$$\text{Mean} = \frac{18}{9} = 2$$

Suppose x_i represents the number whose mean is to be calculated ($x_1, x_2, x_3, x_4, \dots, x_n$). In this set there are n values. So

$$\text{Mean} = \frac{x_1 + x_2 + \dots + x_n}{n} \text{ or } \bar{x} = \frac{\sum x}{n}, \text{ where } \sum x = x_1 + x_2 + x_3 + \dots + x_n.$$

Σ is capital sigma, the Greek letter for S. The notation $\sum x$ stands for “the sum of the x ’s”.

If data is in the form of a frequency distribution, the mean is calculated using the formula:

$$\text{Mean} = \frac{\sum x.f}{\sum f}, \text{ where } \sum x.f \text{ means ‘the sum of the products’}$$

i.e. $\Sigma(\text{number} \times \text{frequency})$ and Σf means ‘the sum of the frequencies’.

Example 4.9

The marks obtained by 100 students in a test were as follows:

Mark (x)	0	1	2	3	4
Frequency (f)	4	19	25	29	23

Find the mean mark.

Solution

Mark (x)	Freq. (f)	x.f
0	4	0
1	19	19
2	25	50
3	29	87
4	23	92
total	100	248

$$\text{Mean} = \frac{248}{100} = 2.48.$$

For grouped data, each class can be represented approximately by its mid-point (class mark)

Example 4.10

The results of 24 students in a Mathematics test are given in the table.

Mark	Freq.	Mid-point(x)	x.f
85 - 99	4	92	368
70 - 84	7	77	539
55 - 69	8	62	496
40 - 54	5	47	235
	$\Sigma f=24$		$\Sigma x.f= 1,638$

$$\text{Mean} = \frac{1,638}{24} = 68.25.$$

Example 4.11.

The number of letters delivered to the 26 houses in a street was as follows:

Number of letters	Number of houses (i.e. freq.)
0 - 2	10
3 - 4	8
5 - 7	5
8 - 12	3

Calculate an estimate of the mean number of letters delivered per house.

Using an assumed mean

When the number of members in a list is large, the above method of finding the mean is quite arduous. From the data we can guess the expected value of the mean. This expected value is called the **assumed** or **working mean**. The assumed mean is approximately in the middle of the data. In a frequency distribution table, the modal value provides the best assumed value. The assumed mean does not need to be one of the values given.

Example 4.12

The table below shows the profit made by a trader in 100 days.

Profit in '000 Ush.	115	125	135	145	155	165
No. of days	8	18	30	26	12	6

Calculate the mean profit

Solution

To calculate the mean of the above data using an assumed mean, the following steps should be followed.

Step 1: Choose an appropriate assumed mean (A) for the range of the values given.

Step 2: Find by how much each of the values (x) differs from this assumed mean (A). These differences obtained are called deviations (d). Thus, $d = x - A$.

Step 3: Multiply each frequency (f) by its corresponding value of deviation (d) to obtain the product (fd).

Step 4: Find the sum of the products, $\sum(fd)$, and then divide this sum by the sum of the frequencies to obtain the mean of the deviations, $\frac{\sum fd}{\sum f}$

Step 5: Find the mean by using the formula: $\bar{x} = A + \frac{\sum fd}{\sum f}$

The table below is a summary of the calculations where the assumed mean is 135 (the modal value).

Profit (x)	Deviations $d = x - A$	Freq. (f)	Products (fd)
115	-20	8	-160
125	-10	18	-180
135	0	30	0
145	10	26	260
155	20	12	240
165	30	6	180
		$\sum f = 100$	$\sum(fd) = 340$

The actual mean = assumed mean + mean deviations

$$= 135 + \left(\frac{340}{100}\right) = 135 + 3.4 = 138.40$$

The mean profit is sh. $138.40 \times 1000 = \text{sh. } 138,400$.

The Median

The median of a set of data is a number selected to represent the middle position when the data are arranged in order of size. The middle position in an

array of N data items is the position numbered $\frac{(N+1)}{2}$. If N is odd, there is a

data item at the middle and we take this item as the median. If N is even, we take the average of the two middle data items as median.

Example 4.13

The median of 18, 22, 30, 31, 44, 60 and 68 is 31 obtained as follows:

Since $N = 7$, position of the median is $\frac{7+1}{2} = 4^{\text{th}}$ data item.

Example 4.14

Find the median of the following numbers. 4, 4, 10, 3, 3, 6, 7, 4, 6, 7.

Solution

Arranging numbers in order of size, we have
3, 3, 4, 4, 4, 6, 6, 7, 7, 10.

Position of median: $\frac{10+1}{2} = 5.5$. So the median lies between the 5th and 6th

data items i.e. between 4 and 6. Hence, the median is $\frac{4+6}{2} = 5$.

Example 4.15

The marks obtained by 100 students in a test were as follows.

Mark (x)	0	1	2	3	4
Frequency (f)	4	19	25	29	23

Find the median mark.

Solution

The median is the number between the 50th and 51st numbers. By inspection, both the 50th and the 51st numbers are 3. Therefore, the median = 3 marks.

It should be noted that, the formula $\frac{(N+1)}{2}$ is not a formula for the median; it simply tells us its position.

For grouped data, the median is given by the following formula:

$$\text{Median} = L_b + \left(\frac{\frac{N}{2} - F_{m-1}}{f_m} \right) \times i.$$

Where, L_b = lower class boundary of the median class;

F_{m-1} = cumulative frequency of the class before the median class;

f_m = frequency of the median class;

i = class width of the median class;

N = total frequency.

Example 4.16

The table below shows the weights of 40 poles in kg.
Find the median weight.

Weight (kg)	frequency	Cumulative freq.
118 - 126	3	3
127 - 135	5	8
136 - 144	9	17
145 - 153	12	29
154 - 162	5	34
163 - 171	4	38
172 - 180	2	40

We find the median class first. Since $\frac{40}{2} = 20$, we look for where 20 is cumulated in the cumulative frequency column. We note that at cumulative frequency of 17, we need 3 more to get 20. So 3 are found in the cumulative of 29. And so '145 - 153' is the median class.


Thus, $L_b = 144.5$; $F_{m-1} = 17$; $f_m = 12$; $i = 9$.

$$\begin{aligned} \text{Hence, median} &= 144.5 + \frac{20 - 17}{12} \times 9 \\ &= 144.5 + 2.25 \\ &= 146.75 \text{ kg.} \end{aligned}$$

Median from an Ogive:

A cumulative frequency curve shows the median at the 50th percentile of the cumulative frequency.

For instance, in figure 5, $N = 62$ and therefore the 31st data item gives the median which is obtained by drawing a horizontal line, from the cumulative frequency axis to the curve, and then drawing the line down wards to the horizontal axis. From the Ogive, the median mark is 49.



The mode

Another measure which is sometimes used to describe the “middle” of a set of data is the mode. It is defined as the value which occurs with the highest frequency.

Example 4.17

Find the mode of the following numbers: 5, 4, 10, 3, 3, 4, 7, 4, 6, 5. The mode is 4. (There are more 4’s than any other number).

For grouped data, the mode is estimated using the following formula.

$$\text{Mode} = l_b + \frac{d_1}{d_1 + d_2} \times i$$

Where, l_b = lower class boundary of the modal class (i.e. the class containing the mode);

d_1 = the difference between the frequency of the modal class and the frequency of the class before it;

d_2 = the difference between the frequency of the modal class and the frequency of the class after it;
 i = class width of the modal class.

Example 4.18

Find the mode of the following data.

Cass	Frequency
20 - 22	3
23 - 25	6
26 - 28	12
29 - 31	9
32 - 34	2

Solution


The modal class is the class with the highest frequency or frequency density (in case the classes have unequal widths).

Thus the modal class for this data is '26 - 28'.

So $l_b = 25.5$, $d_1 = 12 - 6 = 6$, $d_2 = 12 - 9 = 3$, $i = 3$.

$$\text{Mode} = 25.5 + \frac{6}{6+3} \times 3 = 25.5 + \frac{18}{9} = 27.5.$$

When a histogram is drawn, the mode is obtained by joining the ends of the highest rectangle to the opposite ends of the rectangles next to it as shown in the figure below.



Quartiles

These are values that divide the data into four equal parts. They are denoted by Q_1 , Q_2 and Q_3 . Q_1 divides the data into $\frac{1}{4}$, Q_2 into $\frac{2}{4}$ and Q_3 into $\frac{3}{4}$. In other words, for Q_1 , 25% of the values are below it, Q_2 takes 50% and Q_3 takes 75%. Q_1 is called the first (or lower) quartile, Q_2 the second quartile (median) and Q_3 the third or upper quartile.

For **ungrouped data**, the position of Q_1 is given by $\frac{1}{4}(n+1)$ th value, while that of Q_3 by $\frac{3}{4}(n+1)$ th value. In each case n is the number of observations.

Example 4.19

- (i) Find the lower and upper quartiles of the following set of numbers: 3, 12, 4, 6, 8, 5, 4.

Arranging the numbers in order of size, we have
3, 4, 4, 5, 6, 8, 12.

Position of Q_1 : $\frac{1}{4}(7+1)$ th = 2nd value.

Therefore, the lower quartile is 4.

Position of Q_3 : $\frac{3}{4}(7+1)$ th = 6th value.

Therefore, the upper quartile is 8.

For **grouped data**, the following formulae are used:

$$Q_1 = L_b + \left(\frac{\frac{N}{4} - F_{q-1}}{f_q} \right) \times i; \quad Q_3 = L_b + \left(\frac{\frac{3N}{4} - F_{q-1}}{f_q} \right) \times i$$

Where,

L_b = lower class boundary of the quartile class;

F_{q-1} = cumulative frequency of the class before the quartile class;

f_q = frequency of that quartile class;

i = class width of the quartile class;

N = the number of observations in the data.

Example 4.20

The table below shows the distribution of marks gained by a group of students in a mathematics test marked out of 50.

Estimate the lower and upper quartiles.

Marks	Frequency	Cum. Freq.
1 - 10	15	15
11 - 20	20	35
21 - 30	32	67
31 - 40	26	93
41 - 50	7	100

Position of Q_1 is $\frac{1}{4} \times 100 = 25^{\text{th}}$ observation. Using the cumulative frequency column, the 25^{th} observation is located in the '11 - 20' class. This is the lower quartile class.

So, $L_b = 10.5$, $F_{q-1} = 15$, $f_q = 20$, $i = 20.5 - 10.5 = 10$.

$$\begin{aligned} \text{Therefore, } Q_1 &= 10.5 + \left(\frac{\frac{100}{4} - 15}{20} \right) \times 10 \\ &= 10.5 + 5 = 15.5 \text{ marks.} \end{aligned}$$

Position of Q_3 : $\frac{3}{4} \times 100 = 75^{\text{th}}$ observation (mark). Using the cumulative frequency column, the 75^{th} observation is located in the '31 - 40' class.

So, $L_b = 30.5$, $F_{q-1} = 67$, $f_q = 26$, $i = 40.5 - 30.5 = 10$, $N = 100$.

$$\begin{aligned} \therefore Q_3 &= 30.5 + \left(\frac{75 - 67}{26} \right) \times 10 \\ &= 30.5 + 3.0769 = 33.5769 \\ &\cong 33.58 \text{ marks.} \end{aligned}$$

We define the **inter-quartile range** as: $Q_3 - Q_1$,


And **semi-interquartile range** as: $\frac{Q_3 - Q_1}{2}$.

From the Ogive, the value at the 25^{th} percentile is the lower quartile, and that at the 75^{th} percentile is the upper quartile.

Example 4.21

Plot an Ogive for the above data and use it to estimate the semi-interquartile range.

Marks	Frequency	U.C.B	Cum. Freq.
	0	0.5	0
1 - 10	15	10.5	15
11 - 20	20	20.5	35
21 - 30	32	30.5	67
31 - 40	26	40.5	93
41 - 50	7	50.5	100



Example 4.22

The following frequency distribution table gives the marks obtained by 500 candidates in an examination.

Mark	frequency
$1 \leq x < 10$	15
$10 \leq x < 20$	25
$20 \leq x < 30$	39
$30 \leq x < 40$	77
$40 \leq x < 50$	101
$50 \leq x < 60$	84
$60 \leq x < 70$	61
$70 \leq x < 80$	53
$80 \leq x < 90$	22
$90 \leq x < 100$	23


- (a) Draw a cumulative frequency curve for the data.
- (b) From your graph estimate:
- the median mark.
 - the pass mark, if 65% of the candidates passed.
 - the number of candidates who scored less than 45 marks.

Solution

- (a) Construct a cumulative frequency table.

Upper boundary	cf
10	15
20	40
30	79
40	156
50	257
60	341
70	402
80	455
90	477
100	500

Plot the cumulative frequency against the upper class boundaries. The figure below shows the cumulative frequency curve.



- (b) (i) The median mark is located $\frac{1}{2}$ (500) th along the cumulative frequency axis and the corresponding mark is 49.
(ii) If 65% passed, then 35% failed. From the graph, the pass mark is 42 (see the dotted line).
(iii) The number of candidates who scored less than 45 marks is 205.

Exercise 4.1

1. In an experiment, 50 people were asked to guess the weight of a mobile phone in grams. The guesses were as follows:

47 39 21 30 42 35 44 36 19 52
23 32 66 29 5 40 33 11 44 22
27 58 38 37 48 63 23 40 53 24
47 22 44 33 13 59 33 49 57 30
17 45 38 33 25 40 51 56 28 64

Construct a frequency table using intervals 0 - 9, 10 - 19, 20 - 29, etc. Hence draw a cumulative frequency curve and estimate:

- (a) the median weight,
(b) the inter-quartile range,
2. In a competition, 30 children had to pick up as many paper clips as possible in one minute using a pair of tweezers. The results were as follows:
- 3 17 8 11 26 23 18 28 33 38
12 38 22 50 5 35 39 30 31 43
27 34 9 25 39 14 27 16 33 49
- Construct a frequency table using intervals of width 10, starting with 1 - 10. From the frequency table, estimate the
- (i) mean,
(ii) median of the distribution.
3. The mean weight of 8 boys is 55 kg and the mean weight of a group of girls is 52 kg. The mean weight of all the children is 53.2 kg. How many girls are there?
4. A group of 50 people were asked how many books they had read in the previous year; the results are shown in the frequency table below. Calculate the mean number of books read per person.

No of books	0	1	2	3	4	5	6	7	8
Frequency	5	5	6	9	11	7	4	2	1

5. The following tables give the distribution of marks obtained by different classes in various tests. For each table find the mean, median and mode.

(a)

Mark	0	1	2	3	4	5	6
Frequency	3	5	8	9	5	7	3

(b)

Mark	15	16	17	18	19	20
Frequency	1	3	7	1	5	3

Exercise 4.2

1. The table below shows the marks obtained, out of 50, by Form 4E students in a Mathematics test.

Mark (x)	No. of students (f)
1 - 10	9
11 - 20	10
21 - 30	11
31 - 40	8
41 - 50	7

Using an assumed mean, calculate the mean mark to the nearest whole number.

2. The marks obtained by 90 students in an end of term examination are given in the table below.

Mark (%)	No. of students
1 - 10	4
11 - 20	5
21 - 30	10
31 - 40	11
41 - 50	12
51 - 60	15
61 - 70	10
71 - 80	9
81 - 90	8
91 - 100	6

Using 55.5 as the assumed mean, calculate the mean mark.

3. The heights of 50 army recruits were measured and tabulated as shown below.

Height (cm)	No. of recruits
151 - 155	8
156 - 160	16
161 - 165	14
166 - 170	10
171 - 175	2

Using 160 cm as an assumed mean, calculate the mean height.

4. The table below shows the ages of students in a training college.

Age	Frequency
$18 \leq x < 20$	7
$20 \leq x < 22$	10
$22 \leq x < 24$	33
$24 \leq x < 26$	21
$26 \leq x < 28$	14
$28 \leq x < 30$	13
$30 \leq x < 32$	10
$32 \leq x < 34$	4

Calculate the median age.

5. The frequency table below shows the marks scored by 45 students in a test.

Mark	frequency
11 - 15	4
16 - 20	0
21 - 25	5
26 - 30	21
31 - 35	10
36 - 40	3
41 - 50	2

- (a) State the modal class.
 (b) Calculate the median mark.
6. The speeds of public service vehicles during a police check are shown in the table below.

Speed (km/h)	No. of vehicles
31 - 40	5
41 - 50	10
51 - 60	15
61 - 70	30
71 - 80	55
81 - 90	70
91 - 100	15

- (a) Draw a cumulative frequency curve for this data.
- (b) Use your graph to estimate:
- the median speed.
 - the lower and upper quartiles.
 - the percentage of vehicles traveling between 64km/h and 72 km/h.

7. The table below shows the distribution of marks of 81 candidates in a UCE Mathematics examination.

Mark	No. of candidates
$1 < x \leq 10$	1
$10 < x \leq 20$	3
$20 < x \leq 30$	9
$30 < x \leq 40$	11
$40 < x \leq 50$	14
$50 < x \leq 60$	19
$60 < x \leq 70$	11
$70 < x \leq 80$	8
$80 < x \leq 90$	4
$90 < x \leq 100$	1

- (a) Draw a cumulative frequency curve to show the data.
- (b) Use your graph to estimate:
- the median mark.
 - the upper quartile.
 - the number of candidates who scored less than 75 marks.
 - The pass mark, if 60% of the candidates passed.

8. The times taken by a group of students to solve a mathematical problem are given below:

Time(min)	5 - 9	10 - 14	15 - 19	20 - 24	25 - 29	30 - 34
No. of students	5	14	30	17	11	3

- (a) Draw a histogram for the data. Use it to estimate the modal time for solving a problem.
- (b) Calculate the mean time of solving a problem.

9. The ages of 36 students are given below.

13 16 16 15 12 14 13 15 16
16 14 15 12 16 13 14 16 12
15 13 15 16 13 13 15 14 16
13 14 16 15 15 12 14 12 13

Make a frequency table and hence represent the age distribution of the students on a frequency polygon.

5. Probability

Probability is a numerical value given to the chance that a certain event will occur.

Probability that an event occurs = $\frac{\text{number of favourable outcomes}}{\text{total number of outcomes}}$

5.1 Definitions

- (i) An **experiment** is any process by which data is obtained.
- (ii) The results of an experiment are called **outcomes**.
- (iii) A set of all possible outcomes that may occur in a particular experiment is called a **sample space**, usually denoted by **S**. For example,
 - (a) when a coin is tossed: $S = \{H, T\}$
 - (b) when two coins are tossed: $S = \{HT, TH, HH, TT\}$.
 - (c) When three coins are tossed:
 $S = \{HHH, HHT, HTH, THH, TTH, THT, HTT, TTT\}$
 - (d) when a die is thrown: $S = \{1, 2, 3, 4, 5, 6\}$, e.t.c.
- (i) An **event** is a set consisting of possible outcomes of an experiment with desired qualities. It is a subset of a sample space.

5.2 Review of terms used in set theory.

1. Intersection of events.

Consider two events A and B of the sample space S, the intersection of these two events is a set given by $E = A \cap B$, i.e. containing sample points common to both A and B.

2. Union of events, denoted by $A \cup B$, is the set of all sample points in either A or B or both.

3. Compliment of an event.

If A is an event of a sample space S, the compliment of A, denoted by A' , is given by the set containing all sample points in S that are not in A.

4. Mutually exclusive events

If two events A and B are mutually exclusive, then they cannot occur at the same time. In other words $A \cap B = \phi$. For example, selecting 'an even number' or selecting a 'one' from a set of numbers; In tossing a coin, the events 'Head' and 'Tails' are mutually exclusive. If you get a 'Head', you cannot get a 'Tail' in the same toss.

If a trial (experiment) has a set of equally likely possible outcomes S , then the probability of the event E occurring is given by


$$P(E) = \frac{n(E)}{n(S)}$$

5.3 Independent events

Two events are independent if the occurrence of one event is unaffected by the occurrence of the other. E.g. Obtaining a 'head' on one coin, and a 'tail' on the other coin when the coins are tossed at the same time.

5.4 Venn diagrams


Probabilities can be illustrated on a Venn diagram. The rectangle represents the entire sample space, and the circle represents the event A , as shown below.




When dealing with events, the words *and* and *or* are sometimes replaced by the symbols \cap and \cup respectively. The event of A not happening is sometimes written as A' .

Combinations of probabilities can be shown on a Venn diagram, as follows.

A and B , i.e. $A \cap B$. This is the overlap of the regions corresponding to A and B as shown in the figure below.



A or B i.e. $A \cup B$ is the region of points in either the A region or the B region (or both). Note that the word *or* is inclusive. 'A or B' means 'A or B or both', as shown below.



5.5 Basic rules of probability

The following are some of the most basic rules of probability.

1. Probabilities are real numbers on the interval from 0 to 1. i.e. $0 \leq P(E) \leq 1$ for any event E.
2. If an event is certain to occur, its probability is 1, and if an event is certain not to occur, its probability is 0.
3. If A and B are any two events, then

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B).$$
4. If two events are **mutually exclusive**, the probability that one or the other will occur equals the sum of their probabilities
 i.e. $P(A \text{ or } B) = P(A) + P(B)$. This is the addition law for mutually exclusive events. It is also known as the 'or rule'
5. The sum of the probabilities that an event will occur and that it will not occur is equal to 1. i.e. $P(A) + P(A') = 1$ for any event A, or $P(A') = 1 - P(A)$.
6. If A and B are independent events then

$$P(A \text{ and } B) = P(A) \times P(B)$$
 This is the multiplication law for independent events. It is also known as the 'and rule'

In questions on probability, when the following terms are used, great care must be taken. To understand the terms we take, for example a family of 4 children, where B represents a boy and G represents a girl.

- (i) **at least**
 'at least two are boys, $B \geq 2$. Therefore, B can take values 2, 3 or 4.
- (ii) **'at most'**
 'at most two are boys', i.e. $B \leq 2$. Therefore, B can take values 2, 1 or 0.
- (iii) **Not more than**
 'Not more than two boys' and 'at most two boys' mean the same.
- (iv) **Not less than**

'not less than two boys' and 'at least two boys' mean the same.

(v) **More than**

'more than two boys', i.e. $B > 2$. Therefore B can take values 3 or 4.

(vi) **Less than**

'less than two boys', i.e. $B < 2$. Therefore, B can take values 0 or 1.

Example 5.1

The numbers 1 to 20 are each written on a card. The 20 cards are mixed together. One card is chosen at random from the pack. Find the probability that the number on the card is:

(a) even

(b) a factor of 24

(c) prime.

Solution

We will use 'P(x)' to mean 'the probability of x'. Let S be the sample space such that $S = \{1, 2, 3, 4, 5, 6, 7, 8, \dots, 20\}$

$$(a) P(\text{even}) = \frac{\text{number of even numbers}}{\text{total number of numbers in the pack}} = \frac{10}{20} = \frac{1}{2}$$

$$(b) P(\text{a factor of 24}) = \frac{\text{number of factors of 24}}{\text{total number of numbers in the pack}}$$

The factors of 24 are: $F_{24} = \{1, 2, 3, 4, 6, 8, 12, 24\}$.

$$n(F_{24}) = 8$$

$$\text{Therefore, } P(\text{a factor of 24}) = \frac{8}{20} = \frac{2}{5}$$

(c) Prime numbers in the pack = $\{2, 3, 5, 7, 11, 13, 17, 19\}$

$$P(\text{prime}) = \frac{\text{number of prime numbers in the pack}}{\text{total number of numbers in the pack}} = \frac{8}{20} = \frac{2}{5}$$

Example 5.2

A black die and a white die are thrown at the same time. Display all the possible outcomes. Find the probability of obtaining:

(a) a total of 5,

(b) a total of 11,

(c) a 'two' on the black die and a 'six' on the white die.

Solution

It is convenient to display all the possible outcomes on a grid. There are 36 possible outcomes, shown here below.

		Number on white die					
		1	2	3	4	5	6
Number on black die	1	(1, 1)	(1, 2)	(1, 3)	(1, 4)	(1, 5)	(1, 6)
	2	(2, 1)	(2, 2)	(2, 3)	(2, 4)	(2, 5)	(2, 6)
	3	(3, 1)	(3, 2)	(3, 3)	(3, 4)	(3, 5)	(3, 6)
	4	(4, 1)	(4, 2)	(4, 3)	(4, 4)	(4, 5)	(4, 6)
	5	(5, 1)	(5, 2)	(5, 3)	(5, 4)	(5, 5)	(5, 6)
	6	(6, 1)	(6, 2)	(6, 3)	(6, 4)	(6, 5)	(6, 6)

- (a) There are four ways of obtaining a total of 5 on the two dice. These are: (1, 4), (4, 1), (2, 3) and (3, 2)

$$\text{Probability of obtaining a total of 5} = \frac{4}{36} = \frac{1}{9}.$$

- (b) There are two ways of obtaining a total of 11.

$$P(\text{total of 11}) = \frac{2}{36} = \frac{1}{18}$$

- (c) There is only one way of obtaining a 'two' on the black die and a 'six' on the white die.

$$\text{Therefore, } P(\text{a two on black and a 6 on white}) = \frac{1}{36}$$

Example 5.3

A fair coin is tossed and a fair die is rolled. Find the probability of obtaining a 'head' and a 'six'.

Solution

When a coin is tossed once, the sample space is: $S = \{H, T\}$

Where H denotes a 'head' and T a 'tail'. So $P(H) = P(T) = \frac{1}{2}$.

Similarly the sample space when a die is tossed once is: $S = \{1, 2, 3, 4, 5, 6\}$

$$P(\text{six}) = \frac{1}{6}$$

The two events are independent.

$$\text{Therefore, } P(\text{head and six}) = \frac{1}{2} \times \frac{1}{6} = \frac{1}{12}.$$

Exercise 5.1

In this exercise, all dice are normal cubic dice with faces numbered 1 to 6.

- A fair die is thrown once. Find the probability of obtaining:
 - a six,
 - an even number,
 - a number greater than 3,
 - a three or a five.

2. The two sides of a coin are known as 'head' and 'tail'. Two coins are tossed at the same time. List all the possible outcomes. Find the probability of obtaining
 - (a) two heads,
 - (b) a head and a tail.
3. A bag contains 6 red balls and 4 green balls.
 - (a) Find the probability of selecting at random:
 - (i) a red ball
 - (ii) a green ball.
 - (b) One red ball is removed from the bag. Find the new probability of selecting at random:
 - (i) a red ball
 - (ii) a green ball.
4. One letter is selected at random from the word 'UNNECESSARY'. Find the probability of selecting:
 - (a) an R
 - (b) an E
 - (c) an O
5. Three coins are tossed at the same time. List all the possible outcomes. Find the probability of obtaining:
 - (a) three heads,
 - (b) two heads and one tail,
 - (c) no heads,
 - (d) at least one head.
6. A bag contains 10 red balls, 5 blue balls and 7 green balls. Find the probability of selecting at random:
 - (a) a red ball,
 - (b) a green ball,
 - (c) a blue or a red ball,
 - (d) a red or a green ball.
7. A red die and a blue die are thrown at the same time. List all the possible outcomes in a systematic way. Find the probability of obtaining:
 - (a) a total of 10.
 - (b) a total of 12.
 - (c) a total less than 6.
 - (d) the same number on both dice.
 - (e) a total more than 9.
8. A die is thrown; when the result has been recorded, the die is thrown a second time. Display all the possible outcomes of the two throws. Find the probability of obtaining:
 - (a) a total of 4 from the two throws.
 - (b) a total of 8 from the two throws,
 - (c) a total between 5 and 9 inclusive from the two throws,
 - (d) a number on the second throw which is four times the number on the first throw.
9. One ball is selected at random from a bag containing 12 balls of which x

are white.

(a) What is the probability of selecting a white ball?

When a further 6 white balls are added the probability of selecting a white ball is doubled.

(c) Find x .

10. A coin is tossed and a die is thrown. Write down the probability of obtaining:
- (a) a head on the coin,
 - (b) an odd number on the die,
 - (c) a 'head' on the coin and an odd number on the die.
11. A ball is selected from a bag containing 3 red balls, 4 black balls and 5 green balls. The first ball is replaced and a second is selected. Find the probability of obtaining
- (a) two red balls,
 - (b) two green balls.
12. The letters of the word 'INDEPENDENT' are written on individual cards and the cards are put into a box. A card is selected at random and then replaced and then a second card is selected. Find the probability of obtaining:
- (a) the letter 'P' twice,
 - (b) the letter 'E' twice.
13. The table below shows the number of children in 40 randomly chosen families.

No. of children (x)	0	1	2	3	4	5	Over 5	Total
No. of families (f)	6	8	12	5	3	6	0	40

State the probability that a family chosen at random has:

- (a) more than 2 children,
 - (b) an odd number of children,
 - (c) $P(2 \leq x < 4)$.
14. Two tetrahedral dice (4 sided), each with faces labeled 1, 2, 3 and 4 are thrown. The score is the sum of the two numbers on which the dice land. Find the possibility space and the probability of each element of the space.
15. Two dice (6 sided), each with faces labeled 1, 2, 3, 4, 5, 6 are thrown. The score is the difference of the numbers showing on the top faces on two dice. For example, (5, 2) would score 3, as would (2, 5). Find the possibility space and the probability of each score.

TREE DIAGRAMS


These are used to:

- (i) generate a sample space,
- (ii) solve certain probability problems.

Example 5.4

A bag contains 5 red balls and 3 green balls. A ball is drawn at random and then replaced. Another ball is drawn. What is the probability that both balls are green?

Solution



The branch marked * involves the selection of a green ball twice. The probability of this event is obtained by simply multiplying the fractions on the two branches.

$$\text{Therefore, } P(\text{two green balls}) = \frac{3}{8} \times \frac{3}{8} = \frac{9}{64}.$$


Rules:

- (1) When the required outcome is given by a path along the branches of a probability tree, **multiply** the probabilities along that path.
- (2) When the required outcome is given by more than one path in a probability tree, **add** the probabilities resulting from each path.

Example 5.5

A bag contains 5 red balls and 3 green balls. A ball is selected at random and not replaced. A second ball is then selected. Find the probability of selecting:

- (a) two green balls
- (b) one red ball and one green ball.



$$(a) P(\text{two green balls}) = \frac{3}{8} \times \frac{2}{7} = \frac{3}{28}.$$

$$(b) P(\text{one red, one green}) = \left(\frac{5}{8} \times \frac{3}{7}\right) + \left(\frac{3}{8} \times \frac{5}{7}\right) = \frac{15}{28}.$$

Example 5.6

One ball is selected at random from a bag containing 5 red balls, 2 yellow balls and 4 white balls. Find the probability of selecting a red ball or white ball.

Solution

The two events are exclusive.

$$\begin{aligned} P(\text{red ball or white ball}) &= P(\text{red}) + P(\text{white}) \\ &= \frac{5}{11} + \frac{4}{11} = \frac{9}{11}. \end{aligned}$$

Example 5.7

A coin is biased so that it shows 'heads' with a probability of $\frac{2}{3}$. The same coin is tossed three times. Find the probability of obtaining:

- | | |
|---|---------------------|
| (a) two tails on the first two tosses. | Ans. $\frac{1}{9}$ |
| (b) a head, a tail and a head (in that order) | Ans. $\frac{4}{27}$ |
| (c) two heads and one tail (in any order). | Ans. $\frac{4}{9}$ |

Exercise 5.2

1. A bag contains 5 red balls, 3 blue balls and 2 yellow balls. A ball is drawn and not replaced. A second ball is drawn. Find the probability of drawing
- two red balls
 - one blue ball and one yellow ball,
 - two yellow balls,
 - two balls of the same colour.

ANSWERS: (a) $\frac{2}{9}$; (b) $\frac{2}{15}$; (c) $\frac{1}{45}$; (d) $\frac{14}{45}$

2. A six sided die is thrown three times. Draw a tree diagram, showing at each branch the two events: 'six' and 'not six'. What is the probability of throwing a total of
- three sixes
 - no sixes
 - one six
 - at least one six.

ANSWERS: (a) $\frac{1}{216}$; (b) $\frac{125}{216}$; (c) $\frac{25}{72}$; (d) $\frac{91}{216}$.

3. A die has its faces marked 0, 1, 1, 1, 6, and 6. Two of these dice are thrown together and the total score is recorded. Draw a tree diagram.
- How many different totals are possible?
 - What is the probability of obtaining a total of 7?

ANSWERS: (a) 6; (b) $\frac{1}{3}$

4. A coin and a die are tossed. All the possible outcomes are shown in the table

		die					
		1	2	3	4	5	6
coin	H	H1	H2	H3	H4	H5	H6
	T	T1	T2	T3	T4	T5	T6

Find:

- P(a tail and even number)
 - P(head)
 - P(1, 2 or 3)
 - P(head and 6)
 - P(head or 6).
5. Two coins are tossed. Represent the possible outcomes on a probability tree and find the probability that
- both are heads

12. In a super market, it was found that out of 100 apples in a box, 10 were bad and could not be sold. Three apple are picked at random from the box (replacing the previous one each time). What is the probability that:
- all the three are good,
 - none of them is good,
 - 2 or 3 are bad.
13. If two dice are thrown together and their scores recorded as ordered pairs (1, 1), (1, 2), e.t.c, construct a table to show all the possible outcomes. Find the probability that they show:
- the same number,
 - different numbers,
 - 2 as one of the numbers.
14. A bag contains red beads, blue beads and green beads. The probability that a bead drawn at random from the bag will be red is $\frac{3}{5}$ and that it will be blue is $\frac{3}{10}$. What will be the probability that it will be green?
If the bag contains a total of 100 beads, determine the number of beads of each colour.
15. What is the set of possible outcomes if a shilling coin and a die are tossed together? Calculate the probability of a head and a number greater than 3 turning up.
16. What is the probability of getting two tails in three throws of a coin? If the first throw shows a head, what is the probability then?
17. In a class of 30 students, 18 are boys and 12 are girls. A teacher chooses three students at random. Find the probability that:
- all three students are girls,
 - at least one of the three students is a boy?
18. In a shooting practice, two soldiers Ali and Ben fire shots alternatively at a target. Ali fires first. The probabilities of Ali and Ben hitting the target with any shot are $\frac{1}{3}$ and $\frac{2}{5}$ respectively. Using a tree diagram, or otherwise, find the probability that:
- they will both miss the target with their first shot,
 - Ben, in his second round, will be the first to hit the target,
 - Ben will hit the target at least once with his first three shots.
19. A family of three children can be represented by BGB, where B, G and B represents boy first, followed by a girl, followed by another boy. Write down all the eight possible codes fro a family of three children.

Taking $P(B) = P(G) = \frac{1}{2}$, find the probability that a family of three children will contain:

- (a) all three boys,
- (b) at least one girl,
- (c) at most one boy.

20. A bag consists of beads of which 5 are red, 7 are blue and 8 are yellow. Three beads are to be taken, at random. Calculate the probability that the beads will be:

- (a) all yellow,
- (b) all three of the same colour
- (c) all three of different colour,
- (d) all three are not of the same colour.

6. Sets

In Book One, we discussed sets. We said that a set was a collection of objects with something in common. These objects are called elements of the set.

6.1 Symbols:

Below are some of the symbols used in sets:

(a) **Curly brackets:** $\{\}$ this means 'the set of'
E.g. $\{\text{people in this room}\}$ means 'the set of people in this room'

(b) **Members (or elements) of sets.**

$\{\text{Vowels}\} = \{a, e, i, o, u\}$. So a is a member of $\{a, e, i, o, u\}$.

The symbol used is \in which means 'is a member of'. Is r a member of $\{a, e, i, o, u\}$?

No, r is not a member of $\{a, e, i, o, u\}$. Here we write: $r \notin \{a, e, i, o, u\}$.

(d) **Number of members**

Consider $A = \{a, e, i, o, u\}$. A has 5 members. We write: $n(A) = 5$. We can also write: $n\{a, e, i, o, u\} = 5$.

(e) **The empty set**

Let $T = \{\text{people with three eyes}\}$. There are no members in this set. So we call it an empty set. The symbol used for empty set is ϕ or $\{\}$. Thus for the above set we write $T = \{\}$ or Φ .

(f) **Intersection of sets**

Suppose we have two sets A and B . Then the intersection of A and B , written as $A \cap B$, is a set consisting of members that are found in both sets.


Example 6.1

It is given that $A = \{p, t, s, q, r\}$ and $B = \{p, q, u\}$.

$A \cap B = \{p, q\}$.

NOTE:


We can use a Venn diagram to represent the above information:



(g) **Union of sets**

Consider the sets $A = \{p, t, s, q, r\}$; $B = \{p, q, u\}$

The single set consisting of the members of A and the members of B, written as $A \cup B$, is called the union of A and B. The shaded region represents the union of A and B.



The union of A and B contains:


- the members of A only
- the members of $A \cap B$
- the members of B only.

Example 6.2

It is given that $A = \{a, b, c, d, e, f\}$ and $B = \{d, e, f, g, h\}$.

- (a) List the members of $A \cup B$
- (b) State $n(A \cup B)$.

Solution




- (a) $A \cup B = \{a, b, c, d, e, f, g, h\}$
- (b) $n(A \cup B) = 8$.

Note:

The members d, e, and f which are common to A and B are listed only once in the union of A and B.

(h) **Subsets**



All the members of A are also members of B. we say A is a **subset** of B. The symbol for 'is a subset of' is \subset . A is a subset of B is written $A \subset B$. The symbol $\not\subset$ means 'is not a subset of'.

(i) **Universal set (ϵ)**

This is a set containing all members in a described set. For example, $\epsilon = \{\text{counting numbers less than } 10\}$ etc

Suppose $A = \{1,2,3,4\}$; and $B = \{3,4,5,6,7,8,9\}$. Then both A and B are subsets of ϵ .

We can define the **compliment of a set** as that set containing members that are in the universal set but are not found in the set under consideration. For example, the compliment of A, written as A' , is given by


$$A' = \{5, 6,7,8,9\} \text{ and } B' = \{1, 2\}$$

Example 6.3

It is given that $\epsilon = \{1,2,3,4,5,6\}$; $A = \{2,3,5\}$; $B = \{3,4,5\}$. List the members of:

- (i) A' , (ii) B' , (iii) $A' \cap B$, (iv) $A \cup B'$, (v) $(A \cap B)'$, (vi) $A' \cap B'$

Solution




From the above figure

- (i) $A' = \{1, 4, 6\}$
- (ii) $B' = \{1, 2, 6\}$
- (iii) $A' \cap B = \{4\}$
- (iv) $A \cup B' = \{1, 2, 3, 5, 6\}$
- (v) $(A \cap B)' = \{1, 2, 4, 6\}$
- (vi) $A' \cap B' = \{1, 6\}$

Example 6.4

In a school, students must take at least one of these subjects: Maths (M), Physics (P) or Chemistry (C). In a group of 50 students, 7 take all three subjects, 9 take Physics and Chemistry only, 8 take Maths and Physics only and 5 take Maths and Chemistry only. Of these 50 students, x take Maths only, x take Physics only and $x + 3$ take Chemistry only. Draw a Venn diagram, find x , and hence find the number taking Maths.

Solution



$$\begin{aligned}
 \text{From the Venn diagram, } x + 8 + x + 5 + 7 + 9 + (x + 3) &= 50 \\
 3x + 32 &= 50 \\
 3x &= 18 \\
 x &= 6
 \end{aligned}$$

Therefore, the number taking Maths is $6 + 8 + 7 + 5 = 26$.

Note: Entries in the Venn diagram should be written in the following order: $(M \cap P \cap C)$ should be entered first; followed by: $n(M \cap P \text{ only})$, $n(P \cap C \text{ only})$, and $n(M \text{ and } C \text{ only})$; then lastly, $n(M \text{ only})$, $n(P \text{ only})$ and $n(C \text{ only})$.

Example 6.5

The population of a certain town has a choice of three daily newspapers; the New Vision (N), The Monitor (M), and Bukedde (B). 40 read N, 35 read M and 60 read B; 7 read N and M, 10 read M and B and 4 read N and B; 34 read no paper at all. If there are 150 people in the town, find the:

- number of people who read all the three,
- number of people who read only one newspaper.
- number of people who read N and M only.
- probability that a person chosen at random from this town reads two newspapers only.

Solution


The information given can be summarized as follows:

$$n(N) = 40, n(M) = 35, n(B) = 60, n(N \cap M) = 7, n(M \cap B) = 10,$$

$$n(N \cap B) = 4, n(N \cup M \cup B)' = 34 \text{ and } n(\epsilon) = 150.$$

$$\text{Let } n(N \cap M \cap B) = x$$

(a)



Note: $n(N \text{ and } M \text{ only}) = 7 - x$; $n(M \text{ and } B \text{ only}) = 10 - x$; $n(N \text{ and } B \text{ only}) = 4 - x$

$$n(N \text{ only}) = 40 - [(7 - x) + x + (4 - x)] = 40 - [11 - x] = 29 + x.$$

$$n(M \text{ only}) = 35 - [(7 - x) + x + (10 - x)]$$

$$= 35 - [17 - x]$$

$$= 18 + x$$

$$n(B \text{ only}) = 60 - [(4 - x) + x + (10 - x)] = 60 - [14 - x]$$

$$= 46 + x.$$

From the Venn diagram,

$$(29 + x) + (7 - x) + (18 + x) + (10 - x) + (46 + x) + (4 - x) + x + 34 = 150$$

$$48 + x = 150$$

Therefore, $x = 2$.

Hence, the number of people who read all the three is 2.

(b) The number of people who read only one subject is given by:
 $n(\text{N only}) + n(\text{M only}) + n(\text{B only}) = (29 + 2) + (18 + 2) + (46 + 2) = 99.$


(c) the number of people who read N and M only = $7 - 2 = 5.$

(d)
$$P(\text{a person reads 2 papers only}) = \frac{(7 - 2) + (10 - 2) + (4 - 2)}{150}$$
$$= \frac{5 + 8 + 2}{150}$$
$$= \frac{15}{150} \text{ or } \frac{1}{10}$$

Note: If ' $n(N \cap M) = 7$ ', this information **does not** mean 7 read N and M *only*.


Exercise 6

1. In a class of 30 girls, 18 play netball and 14 play hockey, whilst 5 play neither. Find the number who play both netball and hockey.
2. In the Venn diagram $n(A) = 10$, $n(B) = 13$, $n(A \cap B) = x$ and $n(A \cup B) = 18$.



- (a) write in terms of x the number of elements in A but not in B.
 - (b) Write in terms of x the number of elements in B but not in A.
 - (c) Write down an equation in x , hence find the $n(A \cap B)$.
3. The sets A and B intersect such that $n(A \cap B) = 7$, $n(A) = 20$ and $n(B) = 23$. Find $n(A \cup B)$.
 4. Twenty boys in a class all play either football or basketball (or both). If thirteen play football and ten play basketball, how many play both sports?
 5. Of the 53 staff at a school, 36 drink tea, 18 drink coffee and 10 drink neither tea nor coffee. How many drink both tea and coffee?

6. All of 60 different vitamin pills contain at least one of the vitamins A, B and C. Twelve have A only, 7 have B only, and 11 have C only. If 6 have all three vitamins and there are x having both A and B only, B and C only, and A and C only, how many pills contain vitamin A?
7. In a group of 59 people, some are wearing hats, gloves or scarves (or a combination of these), 4 are wearing all three, 7 are wearing just a hat and gloves, 3 are wearing just gloves and a scarf and 9 are wearing just a hat and scarf. The number wearing only a hat or only gloves is x , and the number wearing only a scarf or none of the three items is $(x - 2)$. Find x and hence the number of people wearing a hat.
8. In the Venn diagram, $\mathcal{E} = \{\text{pupils in a class of 15}\}$, $G = \{\text{girls}\}$, $S = \{\text{swimmers}\}$, $F = \{\text{pupils who believe in Father Christmas}\}$. A pupil is chosen at random. Find the probability that the pupil:
- can swim
 - is a girl swimmer,
 - is a boy swimmer who believes in Father Christmas.
- Two pupils are chosen at random. Find the probability that:
- both are boys,
 - neither can swim,
 - both are girl swimmers who believe in Father Christmas.




9. In a class of 30 pupils, all pupils are required to take part in at least two sports chosen from football, gymnastics and tennis. 9 do football and gymnastics; 19 do football and gymnastics; 6 pupils do all three sports. Draw a Venn diagram to show this information. Use your diagram to calculate how many pupils do gymnastics and tennis but not football.
10. The population of a certain town have a choice of three daily newspapers; the Monitor, the New Vision and the Red pepper. 10% take all three

newspapers, 20% take only the Monitor, 20% take both the Monitor and Red pepper but not the New Vision, 10% take only the New Vision. 23% take the New Vision and Monitor, 27% take the Red pepper and New Vision and 5% take only the Red pepper. Draw a Venn diagram to show this information and use it to answer the following questions.

What percentage:

- do not take a newspaper?
- have the Monitor or New Vision?
- have the Monitor or Red pepper?
- have the Red pepper or New Vision but not the Monitor?

11. In the Venn diagram, $\mathcal{E} = \{\text{houses in the street}\}$;
 $C = \{\text{houses with central heating}\}$; $T = \{\text{houses with a colour T.V.}\}$
 $G = \{\text{houses with a garden}\}$



- How many houses have gardens?
- How many houses have a colour T.V and central heating?
- How many houses have a colour T.V and central heating and a garden?
- How many houses have a garden but not a T.V. or a central heating?
- How many houses have a T.V and a garden but not a central heating?
- How many houses are there in the street?

7. Formulae

A formula is a mathematical sentence in which one quantity is expressed in terms of other letters or numbers and letters. For example,

Area of a rectangle = length \times breadth, i.e. $A = lb$;

$C = 2\pi r$ is a formula for finding the circumference, C , of a circle of radius r .

The single letter or quantity on the left is called the **subject of the formula**.

The subject of the formula must occur only once, isolated, in the LHS of the equation.

For example in the formula: $v = u + at$, v is the subject of the formula, but in the

equation: $c = \frac{a(b+c)}{c}$, c is not the subject of the formula (as c appears on both sides, i.e. c is not isolated).

Change of the subject.

Frequently it is convenient to change the subject of a formula. For example, consider the formula $C = 2\pi r$. In this form the subject of the formula is C .

However, if we were to divide both sides of the formula by 2π :

$$\frac{C}{2\pi} = \frac{2\pi r}{2\pi}$$
$$\therefore r = \frac{C}{2\pi}$$

Thus we have changed the subject to r .

Example 7.1

Make I the subject in the formula $T = P + I$.

Solution

$$T = P + I$$

Subtracting P from both sides gives,

$$T - P = I$$

$$\therefore I = T - P.$$

Example 7.2

Make the bold letter the subject in each of the following formulae

(a) $A = \frac{h}{2}(\mathbf{a} + b)$

(b) $x = \sqrt{\mathbf{A}}$

(c) $V = \pi r^2 \mathbf{h}$

(d) $\frac{1}{\mathbf{p}} = \frac{1}{x} + \frac{1}{y}$

Solution

$$(a) A = \frac{h}{2}(a+b)$$

Multiplying both sides by 2

$$2A = h(a+b)$$

Expand the right hand side

$$2A = ha + hb$$

$2A - hb = ha$, subtracting hb from both sides

$$\frac{2A}{h} = a, \text{ dividing by } h.$$

$$\therefore a = \frac{2A}{h}$$

$$(b) x = \sqrt{A}$$

Square both sides to get

$$x^2 = A$$

$$\therefore A = x^2$$

$$(c) V = \pi r^2 h \quad \Leftrightarrow \quad \pi r^2 h = V$$

$$r^2 = \frac{V}{\pi h}$$

$$r = \pm \sqrt{\frac{V}{\pi h}}$$

$$(d) \frac{1}{p} = \frac{1}{x} + \frac{1}{y}$$

Remove fractions first by multiplying by LCM pxy

$$xy = py + px$$

$$xy = p(x+y)$$

$$p(x+y) = xy$$

$$p = \frac{xy}{x+y}$$

In general, to change the subject of a formula, i.e. to express one letter in terms of the other letters:

- (i) square both sides if the letter is under a square root sign.
- (ii) Remove fractions by multiplying all the terms by the LCM of the denominators.
- (iii) Arrange the terms containing the subject (required letter) on one side of the equation.
- (iv) Factorise and divide by the coefficient of the subject (required letter).
- (v) Do not alter capital letters to small letters or vice versa.

Example 7.3

The area and circumference of a circle are given by the formula $A = \pi r^2$ and $C = 2\pi r$ respectively. Show that $C = 2\sqrt{\pi A}$

Solution

$$A = \pi r^2, \text{ hence } r^2 = \frac{A}{\pi} \dots\dots\dots(i)$$

$$\text{Now } C = 2\pi r \text{ or } C^2 = 4\pi^2 r^2 \dots\dots\dots(ii)$$

$$C^2 = 4\pi^2 r^2 = 4\pi^2 \times \frac{A}{\pi} = 4\pi A$$

$$\therefore C = \sqrt{4\pi A} = 2\sqrt{\pi A}$$

Example 7.4

(a) If $a + \frac{bx}{c} = dx$, express x in terms of a , b , c and d .

(b) If $\frac{x+d}{c} = \frac{25d}{x-d}$, obtain x in terms of c and d .

Solution

(a) $a + \frac{bx}{c} = dx$

$$ac + bx = cdx$$

$$ac = cdx - bx = x(cd - b)$$

$$x(cd - b) = ac$$

$$x = \frac{ac}{cd - b}$$

(b) $\frac{x+d}{c} = \frac{25d}{x-d} \Leftrightarrow (x+d)(x-d) = 25cd$

$$\Leftrightarrow x^2 - d^2 = 25cd$$

$$\Leftrightarrow x^2 = 25cd + d^2$$

$$x = \pm \sqrt{25cd + d^2}$$

Exercise 7.1

Make the letters given in brackets the subject of the formulae in the following questions.

1. $v = u + at$ (t)

3. $d = \frac{k}{t} m$ (k)

5. $a = \frac{b+5c}{d}$ (c)

7. $s = \frac{1}{2}(u+v)t$ (v)

9. $s = ut + \frac{1}{2}at^2$ (a)

11. $t = a + \frac{2b}{kt}$ (k)

13. $A = \frac{b+Cx}{x}$ (x)

15. $v^2 = u^2 + 2as$ (v)

17. $V = \frac{1}{2}(a+b)hl$ (b)

19. $V = \frac{1}{3}\pi h^2(3r-h)$ (r)

21. $y = \frac{\sqrt{x+5}}{3}$ (x)

23. $a = b + \sqrt{b^2 + c^2}$ (c)

25. $\frac{y}{x} = \sqrt{py^2 + 1}$ (p)

27. $P = \frac{a\sqrt{x^2 + b^2}}{y}$ (x)

2. $T = 8 + \frac{7}{Q}$ (Q)

4. $F = \frac{9}{5}C + 32$ (C)

6. $R = \frac{p}{Q+5}$ (Q)

8. $I = \frac{PRT}{100}$ (P)

10. $A = P\left(1 + \frac{RT}{100}\right)$ (R)

12. $ax + b = cx + d$ (x)

14. $A = \pi r^2 h$ (r)

16. $A = 2x + \frac{y^2}{7}$ (y)

18. $V = \frac{1}{3}\pi r^2 h + \frac{2}{3}\pi r^3$ (h)

20. $A = b\sqrt{c}$ (c)

22. $\frac{T}{2a} = \sqrt{\frac{b}{d}}$ (d)

24. $K = \sqrt{\frac{PT}{S+T}}$ (T)

26. $P = \frac{Ex}{\sqrt{x^2 + F}}$ (x)

28. $r = \sqrt{\frac{A}{\pi}}$ (A)

Substitution

Example 7.5

- (a) If $x = \frac{ab+c}{d}$, find the value of x when $a = 2$, $b = 3$, $c = 4$ and $d = 10$.
(b) If $v = u + at$, find the value of a when $v = 50$, $u = 20$ and $t = 2$.

Solution

- (a) $x = \frac{ab+c}{d}$. When $a = 2$, $b = 3$, $c = 4$ and $d = 10$,

$$x = \frac{2 \times 3 + 4}{10}, \text{ substituting } a, b, c \text{ and } d \text{ with their given values and using BODMAS.}$$

$$x = \frac{6+4}{10} = \frac{10}{10} = 1.$$

- (b) $v = u + at$ when $v = 50$, $u = 20$, $t = 2$,
 $50 = 20 + 2a$ substituting v , u , and t with their given values.

$$50 - 20 = 2a$$

$$2a = 30$$

$$a = \frac{30}{2}$$

$$\therefore a = 15.$$

Exercise 7.2

- If $x = 2ab$, find the value of x when $a = 4$, $b = 5$.
- If $x = \frac{a+b}{c}$, find the value of x when $a = -1$, $b = 9$ and $c = 5$.
- If $b - ax = c$
 - make x the subject
 - find the value of x when
 - $a = 2$, $b = 10$, $c = 8$
 - $a = 3$, $b = 6$, $c = 15$.
- If $A = \frac{1}{2}bh$, find the value of
 - A when $b = 6$ and $h = 5$
 - h when $A = 15$ and $b = 6$
 - b when $A = 10$ and $h = 4$
- If $I = \frac{nrP}{100}$, find the value of

- (a) I when $P = 200$, $r = 5$ and $n = 4$
 (b) P when $I = 15$, $r = 10$ and $n = 3$.
 (c) r when $I = 4.5$, $P = 150$ and $n = 1$.
6. A temperature measured in degrees Fahrenheit (F) can be converted to degrees Celsius (C) using the formula $C = \frac{5}{9}(F - 32)$. Given that on a particular day, the maximum temperature in London was 80°F and in Paris was 30°C , which city had the higher temperature?
7. The distance d km which a person can see when at height h m above the ground is given by the formula $d = 8\sqrt{\frac{h}{5}}$. Find
 (a) the distance seen if the person is 45 m above ground level,
 (b) how high the person must be to see a distance of 16 km.
8. The time (Tsec) taken for one complete swing of a pendulum is related to the length of the pendulum (I m) by the formula

$$T = 2\pi\sqrt{\frac{I}{g}}$$
 where $\pi = 3.14$ and $g = 9.81$ m/s, both correct to 3 significant figures. Find
 (a) the time for one complete swing of a pendulum which is 1 m long.
 (b) the length of pendulum which will result in a complete swing lasting exactly one second.
9. $\frac{1}{u} + \frac{1}{v} = \frac{1}{f}$ is the formula for the focal length of a lens.
 (a) Make f the subject of the formula.
 (b) Find f when $u = \frac{1}{2}$ and $v = \frac{1}{3}$.
10. Make k the subject of the formula: $\frac{1}{n^2} = \frac{k^2 + a^2}{hg}$. Hence, evaluate k if $h = 2$, $n = 1.6$, $a = 3$ and $g = 32$.
11. Make C the subject of the formula $C^2d + 2 = p^2q^2$ and evaluate C when $p = 5$, $q = 2$ and $d = 1.62$.
12. The area A of a trapezium, whose parallel sides 'a' cm and 'b' cm are 'h' cm apart, is given by $A = \frac{h(a+b)}{2}$.
 Make h the subject of the formula.
 Evaluate h if $A = 1.12 \times 10^3$ mm² and the sum of the parallel sides is 3.5×10^2 mm.

13. If $A = \frac{c}{\sqrt{(c-b)(c+b)}}$, make b the subject of the formula. Evaluate A^2 if $c = 2b$.
14. A runner covers a distance d km in t hours. Obtain and simplify an expression for the time T he would take to cover the same distance if he increased his speed by 1 km/h.
Make d the subject of the expression you obtain.
15. A grocer bought butter at sh. A per 100 kg and sold it at sh. B per kg thereby gaining P per cent on the cost price.
Show that $P = \frac{100(5b - a)}{a}$.
Find also a formula giving a in terms of P and b .
16. A positive number is given by the formula $N = \frac{3k}{\sqrt{m}}$
- Express m in terms of N and K .
 - If K may take any value from 5 to 8 (inclusive) and m may take any value from 100 to 225 (inclusive), calculate the greatest value of N .
17. Make M the subject of the formula: $N = 2\pi\sqrt{LM}$.
Evaluate M , correct to 2 decimal places, if $L = 0.117$, $\pi = 3.14$ and $N = 3.55$.
18. Make b the subject of the formula: $d = \sqrt{\left(\frac{a}{b} - c\right)}$.
Evaluate b if $a = 4.52 \times 10^5$, $c = 1.3 \times 10^4$ and $d = 1.87 \times 10^2$
19. The radius R cm of a solid sphere of mass W g and density d g/cm³ is given by the formula:
- $$R = \sqrt[3]{\frac{3W}{4\pi d}}$$
- Make W the subject of the formula. Evaluate W if $R = 0.3$ cm, and $d = 5.316$ g/cm³, giving your answer correct to 3 SF. (Take $\pi = 3.142$).
20. If $\frac{Q^2}{P} = \frac{1}{T^2}$, make T the subject of the formula.
Evaluate T if $P = 1.44 \times 10^{-6}$ and $Q = 4.8 \times 10^{-6}$, giving your answer in standard form.
21. From the formula $(p + \frac{a}{b^2})(b - c) = d$, express p in terms of a , b , c and d without simplifying your answer.
If $a = 8.0 \times 10^{-3}$, $b = 1.2 \times 10^{-2}$, $c = 2.0 \times 10^{-3}$ and $d = 1$, evaluate p .

8. Functions

A function is a rule which associates each element of one set with an element of another.

The first set is called the **domain** and the second set is called the **range**.

The two common notations used are:

(a) $f(x) = x^2 + 4$

(b) $f : x \mapsto x^2 + 4$

We may interpret (b) as follows 'function f is such that x is mapped onto $x^2 + 4$.

Example 8.1


Given domain $A = \{1, 2, 3\}$, range $B = \{1, 2, 3, 4, 5\}$ and function $f : x \mapsto x + 2$, draw a line diagram to illustrate the function.

Solution

$1 \in A$ is associated with $1 + 2 = 3 \in B$

$2 \in A$ is associated with $2 + 2 = 4 \in B$

$3 \in A$ is associated with $3 + 2 = 5 \in B$.




Example 8.2

For each of the following functions, write down the images of -3, 0 and 3.

(a) $f : x \mapsto 2x + 3$

(b) $f : x \mapsto x^2$

Solution




Therefore, the range is $\{-3, 3, 9\}$

(b) The images of $\{-3, 0, 3\}$ under the function $f : x \mapsto x^2$, are $\{9, 0, 9\}$

Example 8.3

Given domain $S = \{2, 3, 4, 5\}$ write down the range of $f : x \mapsto x^2 + 3$

Solution



The range of S is $\{7, 12, 19, 28\}$.

Example 8.4

If $f(x) = 3x - 1$ and $g(x) = 1 - x^2$ find:

- | | |
|-----------------------|-------------|
| (a) $f(2)$ | (b) $f(-2)$ |
| (c) $g(0)$ | (d) $g(3)$ |
| (e) x if $f(x) = 1$ | |

Solution

(a) $f(2) = 3(2) - 1 = 6 - 1 = 5.$

(b) $f(-2) = 3(-2) - 1$
 $= -6 - 1$
 $= -7.$

(c) $g(0) = 1 - 0 = 1.$

(d) $g(3) = 1 - 9 = -8$

(e) If $f(x) = 1$ then $3x - 1 = 1$
 $3x = 2$
 $x = \frac{2}{3}.$

Example 8.5

The function f is defined as $f(x) = ax + b$ where a and b are constants. If $f(1) = 8$ and $f(4) = 17$, find the values of a and b .

Solutions

$f(x) = ax + b$



If $f(1) = 8$, $\Rightarrow a(1) + b = 8$
 $a + b = 8$ (i)
 $f(4) = 17 \Rightarrow 17a + b = 17$(ii)

equation (ii) - equation (i), gives

$$16a = 9$$

$$\Rightarrow a = \frac{9}{16}$$

Substituting for a into equation (i) gives

$$\frac{9}{16} + b = 8$$

$$b = 8 - \frac{9}{16} = \frac{128}{16} - \frac{9}{16} = \frac{119}{16}$$

Therefore, $a = \frac{9}{16}$ and $b = \frac{119}{16}$.

Exercise 8.1

1. Given the functions $h : x \mapsto x^2 + 1$ and $g : x \mapsto 10x + 1$. Find:
 (a) $h(2), h(-3), h(0)$ (b) $g(2), g(10), g(-3)$.

In questions 2, 3 and 4, the functions f, g and h are defined as follows:

$$f : x \mapsto 1 - 2x; \quad g : x \mapsto \frac{x^3}{10}; \quad h : x \mapsto \frac{12}{x}$$



2. Find:
 (a) $f(5), f(-5), f(\frac{1}{4})$
 (b) $g(2), g(-3), g(\frac{1}{2})$
 (c) $h(3), h(10), h(\frac{1}{3})$
3. Find:
 (a) x if $f(x) = 1$ (b) x if $fx) = -11$
 (c) x if $h(x) = 1$
4. Find:
 (a) y if $g(y) = 100$ (b) z if $h(z) = 24$

For questions 5 and 6, the functions k, l and m are defined as follows:

$$k(x) = \frac{2x^2}{3}, \quad l(x) = \sqrt{[(y-1)(y-2)]}, \quad m(x) = 10 - x^2$$

5. (a) $k(3), k(6), k(-3)$ (b) $l(2), l(0), l(4)$
 (c) $m(4), m(-2), m(\frac{1}{2})$
6. Find:
 (a) x if $k(x) = 6$ (b) x if $m(x) = 1$





So $f^{-1}(x) = \frac{x+4}{3}$.

Example 8.7

Find the inverse of the function $f(x) = \frac{3(x-1)}{4}$

Solution

Let $y = \frac{3(x-1)}{4}$

$4y = 3(x-1)$ (multiplying by 4 on both sides)

$4y = 3x - 3$

$4y + 3 = 3x$ Adding 3

$3x = 4y + 3$

$x = \frac{4y+3}{3}$

Replacing y with x in the function $\frac{4y+3}{3}$ we get $\frac{4x+3}{3}$.

Therefore, $f^{-1}(x) = \frac{4x+3}{3}$.

Composite functions

The function $f(x) = 3x + 2$ is itself a composite function, consisting of two simpler functions: 'multiply by 3' and 'add 2'.

If $f(x) = 3x + 2$ and $g(x) = x^2$ then fg is a composite function where g is performed first and then f is performed on the result of g .

Note: $fg \neq gf$ as a general rule. That is, composition of functions is not commutative.

Example 8.8

Given that: $f(x) = 3x + 2$ and $g(x) = x^2$. Find

- | | |
|-----------------|--------------------|
| (a) $fg(x)$ | (b) $gf(x)$ |
| (c) f^{-1} | (d) g^{-1} |
| (e) $(gf)^{-1}$ | (f) $g^{-1}f^{-1}$ |

Solution

$$\begin{aligned} \text{(a) } fg(x) &= f[g(x)] \\ &= f[x^2] = 3(x^2) + 2 \\ &= 3x^2 + 2 \end{aligned}$$

$$\begin{aligned} \text{(b) } gf(x) &= g[f(x)] \\ &= g[3x + 2] \\ &= (3x + 2)^2 \end{aligned}$$

$$\begin{aligned} \text{(c) } f(x) &= 3x + 2 \\ \text{Let } y &= 3x + 2 \\ y - 2 &= 3x \quad \text{or} \quad 3x = y - 2 \\ &\therefore x = \frac{y - 2}{3} \\ &\therefore f^{-1}(x) = \frac{x - 2}{3}. \end{aligned}$$

$$\text{(d) } g(x) = x^2 \Rightarrow g^{-1}(x) = \sqrt{x}.$$

$$\begin{aligned} \text{(e) } gf(x) &= (3x + 2)^2 \\ \text{Let } y &= (3x + 2)^2 \\ \sqrt{y} &= 3x + 2 \Rightarrow 3x + 2 = \sqrt{y} \\ 3x &= \sqrt{y} - 2 \\ \Rightarrow x &= (\sqrt{y} - 2)/3 \end{aligned}$$

$$\text{Hence, } (gf)^{-1} = \frac{\sqrt{x} - 2}{3}$$

$$\text{(f) } g^{-1}f^{-1} = g^{-1}[f^{-1}(x)] = g^{-1}\left[\frac{x - 2}{3}\right]$$

$$\text{But } g^{-1}(x) = \sqrt{x} \Rightarrow g^{-1}\left[\frac{x - 2}{3}\right] = \sqrt{\frac{x - 2}{3}}$$

$$\text{Therefore, } g^{-1}f^{-1} = \sqrt{\frac{x - 2}{3}}.$$

Rational functions

If $h(x) = \frac{f(x)}{g(x)}$, then $h(x)$ is called a rational function.

The function $\frac{f(x)}{g(x)}$ becomes **equal to zero** when the numerator is equal to zero, i.e. when $f(x) = 0$.

The function $\frac{f(x)}{g(x)}$ is **undefined** or **meaningless** when the denominator is equal to zero, i.e. when $g(x) = 0$

Example 8.9

Given that $h(x) = \frac{x+7}{x^2+3x+2}$. Find the value(s) of x for which

- (a) $h(x) = 0$
- (b) $h(x)$ if undefined.

Solution

$$\begin{aligned} \text{(a) } h(x) = 0 &\Rightarrow \frac{x+7}{x^2+3x+2} = 0 \\ &\Rightarrow x+7 = 0 \\ &\Rightarrow x = -7 \end{aligned}$$

$$\begin{aligned} \text{(b) } h(x) \text{ is undefined when } x^2+3x+2 &= 0 \\ (x+2)(x+1) &= 0 \\ x+2 = 0 &\Rightarrow x = -2 \\ \text{Or } x+1 = 0 &\Rightarrow x = -1 \end{aligned}$$

Therefore, $x = -2$ or $x = -1$.

Exercise 8.2

Find the inverse of each the following functions.

1. $f(x) = x + 3$

2. $f(x) = 2x$

3. $f(x) = 3x - 1$

4. $2x + 3$

5. $f(x) = 7x - 3$

6. $h(x) = \frac{1}{2}x + 6$

7. $g(x) = x^3$

8. $f(x) = x^3 + 7$

9. $f : x \rightarrow \frac{2x+3}{5}$

10. $f : x \rightarrow \frac{1}{x+1}$

Exercise 8.3

For questions 1 and 2, the functions f , g and h are as follows:

$$f(x) = 4x, \quad g(x) = x + 5, \quad h(x) = x^2$$

1. Find the following

- | | |
|--------------|--------------|
| (a) $fg(x)$ | (b) $gf(x)$ |
| (c) $hf(x)$ | (d) $fh(x)$ |
| (e) $gh(x)$ | (f) $fgh(x)$ |
| (g) $hfg(x)$ | |

2. Find:

- | | |
|---------------------------|----------------------------|
| (a) x if $hg(x) = h(x)$ | (b) x if $fh(x) = gh(x)$ |
|---------------------------|----------------------------|

For questions 3, 4 and 5, the functions f , g and h are as follows:

$$f(x) = 2x, \quad g(x) = x - 3, \quad h(x) = x^2$$

3. Find the following composite functions

- | | |
|-----------|-----------|
| (a) fg | (b) gf |
| (c) gh | (d) hf |
| (e) ghf | (f) hgf |

4. Evaluate:

- | | |
|---------------|---------------|
| (a) $fg(4)$ | (b) $gf(7)$ |
| (c) $gh(-3)$ | (d) $fgf(2)$ |
| (e) $ggg(10)$ | (f) $hfh(-2)$ |

5. Find:

- | | |
|--------------------------|----------------------------|
| (a) x if $f(x) = g(x)$ | (b) x if $hg(x) = gh(x)$ |
| (c) x if $gf(x) = 0$ | (d) x if $fg(x) = 4$ |
| (e) $h^{-1}g^{-1}(x)$ | (f) $(fgh)^{-1}$ |

Find the value(s) of x for which each of the following functions is:

- | | |
|-----------------|------------------|
| (a) equal to 0, | (b) meaningless. |
|-----------------|------------------|

6. $f(x) = \frac{x+2}{x^2+7x+12}$

7. $f(x) = \frac{x^2-1}{2x^2+7x-15}$

8. $g(x) = \frac{9x^2-1}{x^2+5x-14}$


9. $h(x) = \frac{x(x+5)}{2x+1}$

10. $f(x) = \frac{x+1}{3x+2}$

11. $f(x) = \frac{1+4x}{8(x+2)}$

9. Trigonometry (2)

Triangle PQR in the figure below is right-angled and $\angle PQR = \theta$.



In the figure, PQ is the hypotenuse; PR is the opposite side to θ .
In general:

$$\sin \theta = \frac{\text{opposite side}}{\text{hypotenuse}} = \frac{PR}{PQ}$$

$$\cos \theta = \frac{\text{adjacent side}}{\text{hypotenuse}} = \frac{QR}{PQ}$$


$$\tan \theta = \frac{\text{opposite side}}{\text{adjacent side}} = \frac{PR}{QR}$$

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

Angles of 45° , 60° and 30°

Tan, sin and cos of 45°

Consider an isosceles right-angled triangle ABC such that $AB = BC = 1$ unit and $\angle B = 90^\circ$



$$AC^2 = 1^2 + 1^2 = 2 \quad (\text{Pythagoras' theorem})$$


Therefore, $AC = \sqrt{2}$.

Since $AB = BC$, $\angle A = \angle C = 45^\circ$

$$\tan 45^\circ = \frac{1}{1} = 1; \quad \sin 45^\circ = \frac{1}{\sqrt{2}}; \quad \cos 45^\circ = \frac{1}{\sqrt{2}}$$

Tan, sin and cos of 60° and 30°

Consider an equilateral triangle ABC whose sides are 2 cm as shown in the diagram below. AN is its perpendicular height.



Using Pythagoras' theorem, we have

$$BN^2 + AN^2 = 2^2$$

$$1 + AN^2 = 4$$

$$AN^2 = 3$$

Therefore, $AN = \sqrt{3}$.

$$\sin 60^\circ = \frac{\sqrt{3}}{2}; \quad \cos 60^\circ = \frac{1}{2} \quad \text{and} \quad \tan 60^\circ = \frac{\sqrt{3}}{1} = \sqrt{3}.$$

$$\sin 30^\circ = \frac{1}{2}; \quad \cos 30^\circ = \frac{\sqrt{3}}{2} \quad \text{and} \quad \tan 30^\circ = \frac{1}{\sqrt{3}}$$

Example 9.1

If $\cos \theta = \frac{1}{\sqrt{2}}$, where θ is acute, without using tables or calculator, find:

(a) $\sin \theta$ (b) $\tan \theta$.

(c) $\cos \theta + \tan \theta$. Leave your answers in surd form.

Solution


(a) Consider $\triangle ABC$ right-angled at B. Using Pythagoras' theorem,

$$AB^2 + BC^2 = AC^2$$

$$AB^2 + 1^2 = (\sqrt{2})^2$$

$$AB^2 = 2 - 1 = 1$$

Therefore, $AB = 1$.



(a) $\sin \theta = \frac{1}{\sqrt{2}}$ and

(b) $\tan \theta = \frac{1}{1} = 1.$

(c) $\cos \theta + \tan \theta = \frac{1}{\sqrt{2}} + 1$
 $= \frac{1 + \sqrt{2}}{\sqrt{2}} = \frac{\sqrt{2}(1 + \sqrt{2})}{2}$ or $\frac{\sqrt{2} + 2}{2}$


The graph of $y = \sin x$ (Sine Curve)

To draw the graph of $y = \sin x$, first, prepare a table of values of x and y . Choose suitable values of x in the interval $0 \leq x \leq 720^\circ$ and then find the corresponding values of y . The table below shows the values of $y = \sin x$ in the interval $0 \leq x \leq 720^\circ$.

x	0	90	180	270	360	450	540	630	720
$y = \sin x$	0	1	0	-1	0	1	0	-1	0

Plot the ordered pairs (x, y) on a graph paper and join the points with a smooth curve.

The diagram below shows the graph of $y = \sin x$, in the interval $0 \leq x \leq 720^\circ$



The graph of sine curve, $y = \sin x$, shown above illustrates that:


- (i) The sine curve is a wave which lies between -1 and $+1$; it is zero at multiples of 180° , $-1 \leq \sin x \leq 1$
- (ii) The graph repeats itself at intervals of 360° .
- (iii) For any value of $\sin x$ in the range -1 to $+1$, there are two corresponding values of x in the domain $0 \leq x \leq 360^\circ$. (Except when $\sin x = -1, 0$ and 1).

The graph of $y = \cos x$

The table below shows the values of $y = \cos x$ in the interval $0 \leq x \leq 360^\circ$

x	0	60	90	120	180	240	270	300	360
$y = \cos x$	1	0.5	0	-0.5	-1	-0.5	0	0.5	1

Plot the ordered pairs $(x, \cos x)$ on the graph and join them to obtain the required curve as shown below.



The sine rule

The sine rule enables us to calculate sides and angles in some triangles where there is not a right angle. It is used when:


- (i) one side and two angles are known or
- (ii) two sides and an angle (opposite to one of the two sides) are known.

In $\triangle ABC$, we use the convention that

a is the side opposite \hat{A}

b is the side opposite \hat{B}

c is the side opposite \hat{C}



In any triangle ABC, the sine rule states that


$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} \quad \dots\dots\dots (i)$$

Or $\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c} \quad \dots\dots\dots (ii)$

Use (i) when finding a side, and (ii) when finding an angle.

Example 9.2


Find the value of c in triangle ABC



$$\frac{c}{\sin C} = \frac{b}{\sin B}$$

$$\frac{c}{\sin 50^\circ} = \frac{7}{\sin 60^\circ} \Leftrightarrow c = \frac{7 \times \sin 50^\circ}{\sin 60^\circ} = 6.19 \text{ cm (3 s.f.)}$$

Example 9.3



Find \hat{B} .

Solution

$$\frac{\sin B}{b} = \frac{\sin A}{a} \Leftrightarrow \frac{\sin B}{6} = \frac{\sin 120^\circ}{15}$$


$$\sin B = \frac{6 \times \sin 120^\circ}{15} = 0.346$$

$$\hat{B} = 20.3^\circ$$


Exercise 9.1

For questions 1 to 6, find each side marked with a letter. Give answers to 3 S.F.


1.




2.




3.




4.



5.



6.




7. In triangle ABC, $\hat{A} = 61^\circ$, $\hat{B} = 47^\circ$, $AC = 7.2$ cm. Find BC.
8. In $\triangle XYZ$, $\hat{Z} = 32^\circ$, $\hat{Y} = 78^\circ$, $XY = 5.4$ cm. Find XZ.
9. In $\triangle PQR$, $\hat{Q} = 100^\circ$, $\hat{R} = 21^\circ$, $PQ = 3.1$ cm. Find PR.
10. In $\triangle LMN$, $\hat{L} = 21^\circ$, $\hat{N} = 30^\circ$, $MN = 7$ cm. Find LN.
11. In $\triangle ABC$, $\hat{A} = 62^\circ$, $BC = 8$ cm, $AB = 7$ cm. Find \hat{C} .
12. In $\triangle XYZ$, $\hat{Y} = 97.3^\circ$, $XZ = 22$, $XY = 14$. Find \hat{Z} .
13. In $\triangle DEF$, $\hat{D} = 58^\circ$, $EF = 7.2$, $DE = 5.4$. Find \hat{F} .
14. In $\triangle LMN$, $\hat{M} = 127.1^\circ$, $LN = 11.2$, $LM = 7.3$. Find \hat{L} .

The Cosine Rule

We use the cosine rule when we have either

- (a) two sides and the included angle or
- (b) all three sides.



There are two forms:

1. To find the length of a side:

$$a^2 = b^2 + c^2 - 2bc \cos A$$


$$\text{or } b^2 = a^2 + c^2 - 2ac \cos B$$

$$\text{or } c^2 = a^2 + b^2 - 2ab \cos C$$

2. To find an angle when given all three sides.

$$\cos A = \frac{b^2 + c^2 - a^2}{2bc} \quad \text{or} \quad \cos B = \frac{a^2 + c^2 - b^2}{2ac} \quad \text{or} \quad \cos C = \frac{a^2 + b^2 - c^2}{2ab}$$

Example 9.4




Find b.

Solution

$$\begin{aligned} b^2 &= a^2 + c^2 - 2ac \cos B \\ &= 64 + 25 - (2 \times 8 \times 5 \times \cos 112^\circ) \\ &= 64 + 25 - (80 \times (-0.3746)) = 64 + 25 + 29.968 \\ b &= \sqrt{(118.968)} = 10.9 \text{ cm (3 s.f.)} \end{aligned}$$

Example 9.5




Find angle C.

Solution

$$\begin{aligned} \cos C &= \frac{a^2 + b^2 - c^2}{2ab} = \frac{5^2 + 6^2 - 7^2}{2 \times 5 \times 6} = \frac{12}{60} = 0.200 \\ \hat{C} &= 78.5^\circ \end{aligned}$$

The area of a triangle




For any triangle,

Area = half the product of any two sides and the sine of the included angle.

$$= \frac{1}{2}ac \sin B \text{ or } \frac{1}{2}bc \sin A \text{ or } \frac{1}{2}ab \sin C$$

Example 9.6

Find the area of a triangle ABC in which $a = 4.2$ cm, $c = 7.5$ cm, and $B = 110^\circ$.




$$\begin{aligned} \text{Area} &= \frac{1}{2}ac \sin B \\ &= \frac{1}{2} \times 4.2 \times 7.5 \times \sin 110^\circ \\ &= 2.1 \times 7.5 \times \sin 110^\circ \\ &= 14.8 \text{ cm}^2. \end{aligned}$$


Exercise 9.2

In questions 1 - 3, find the length of the sides marked with letters.


1.



2.



3.




In questions 4 - 6, find to the nearest tenth of a degree the size of the marked angles.

4.


5.

6.




Using the Cosine Rule (and Sine Rule if necessary), work out Q.7 - 12 for $\triangle ABC$.

7. $b = 3$ cm, $c = 4.8$ cm and $A = 120^\circ$. Find a and B .
8. $a = 4$ cm, $b = 7$ cm and $C = 42.5^\circ$. Find C , A and B .
9. $a = 7$ cm, $b = 8$ cm and $c = 6$ cm. Find the angles of the triangles
10. $a = 4$ cm, $b = 5$ cm and $C = 120^\circ$. Find c , A and B .
11. $b = 10$ cm, $c = 12$ cm and $c = 20$ cm. Find angle C
12. $b = 10$ cm, $c = 12$ cm and $A = 130^\circ$. Find a , B and C
13. In $\triangle PQR$, $q = 3$ cm, $r = 5$ cm and $P = 120^\circ$. Find p and also the area of the triangle.
14. In $\triangle ABC$, $b = 3$ cm, $c = \frac{1}{2}b$ and $A = 50^\circ$. Find a , C and B .
15. In $\triangle ABC$, $a = 5$ cm, $b = 7$ cm and $c = 9$ cm. Calculate the angle B and the area of the triangle.
16. In $\triangle ABC$, $AB = 3$ cm, $BC = 4$ cm and angle $B = 40^\circ$. Calculate the area of the triangle and the length of the third side.
17. In $\triangle ABC$, $a = 3$ cm, $b = 5$ cm and angle $A = 20^\circ$. Calculate the two possible values of the angle B .
18. The diagonals of a parallelogram $ABCD$ intersect at O , where angle $AOB = 45^\circ$. Find the area of the parallelogram, if the diagonal $AC = 7.2$ cm and the diagonal $BD = 5.6$ cm.
19. In the figure below, $AB = 8$ cm, $BD = 20$ cm, $DC = 14$ cm, angle $BDC = 90^\circ$ and angle $ABC = 76^\circ$.
Calculate:
 - (a) the size of the angle ABD ,
 - (b) area of the quadrilateral $ABCD$.




Exercise 9.3


- PQRS is a trapezium in which PQ is parallel to SR. The diagonal SQ = 5 cm, RS = 7 cm and angle RSQ = 26° . Calculate the length of QR and the area of triangle QSR. Given also that PS = QR, calculate angle SPQ.
- A point X is 10 nautical miles due north of Y. The bearing of a ship S from X is 140° and that from Y is 080° . Calculate the distance of the ship from X.
- ABC is a triangle whose base BC = 35 cm. The point X on BC is such that BX = 21 cm, AX = 16 cm and angle AXB = 60° . Calculate:
 - the length of AB,
 - the length of AC,
 - the size of angle BAC.
- The point P is 5 km due north of the point Q. A man walks from Q in a direction 030° . Calculate how far he walks before he is
 - equidistant from P and Q,
 - as close as possible to P and
 - north east of P.
- P is a point on the coast 1.5 km due east of a guard station C. S is another point on the coast 2.7 km from C. A ship sailing from P to S is intercepted at Q by a boat from the coast guard station. Given that angle PCS = 63° and QS = 0.5 km, calculate:
 - PQ,
 - the bearing of Q from P.



6. A destroyer D and a cruiser C leave port P at the same time. The destroyer sails 25 km on a bearing 040° and the cruiser sails 30 km on a bearing of 320° . How far apart are the ships?
7. Find all the angles of a triangle in which the sides are in the ratio 5 : 6 : 8
8. From A, B lies 11 km away on a bearing of 041° and C lies 8 km away on a bearing of 341° . Find:
 - (a) the distance between B and C
 - (b) the bearing of B from C.
9. From a lighthouse L an aircraft carrier A is 15 km away on a bearing of 112° and a submarine S is 26 km away on a bearing of 200° . Find:
 - (a) the distance between A and S,
 - (b) the bearing of A from S.
10. If the line BCD is horizontal find:
 - (a) AE
 - (b) \hat{EAC}
 - (c) the angle of elevation of E from A.



11. An aircraft flies from its base 200 km on a bearing 162° , then 350 km on a bearing 260° , and then returns directly to base. Calculate the length and bearing of the return journey.
12. Town Y is 9 km due north of town Z. Town X is 8 km from Y, 5 km from Z and somewhere to the west of the line YZ.
 - (a) Draw triangle XYZ and find angle \hat{YZX} .
 - (b) During an earthquake, town X moves due South until it is due West of Z. Find how far it has moved.
- 13.




Calculate WX, given $YZ = 15$ m.

10. Three dimensional geometry


A line has one dimension because to locate a point on it, we move in one direction.



A plane has two dimensions because to locate a point on a plane we use two coordinates (x, y) .




A solid has three dimensions because a point on it is located using three coordinates (x, y, z).




Geometrical properties of solids

Some of the common regular solids are cubes, cuboids, cylinders, pyramids, prisms, cones and spheres.




cube



cuboid



cylinder




Solids may have faces, edges, and vertices. These faces, edges and vertices determine the geometrical properties of a solid. Solids with flat faces are called **Polyhedra** (singular, polyhedron). Polyhedron means many faces. A regular polyhedron consists of faces that are made of regular polygons.

Projection of a line on a plane

In the figure on the next page, AB is a line meeting plane PQRS at point B. The line is inclined to the plane at angle θ .


Line AC is perpendicular to the plane from point A. Line BC is referred to as the **projection** of line AB on plane PQRS. The projection of line AB on plane PQRS is the shadow of line AB when the source of light is perpendicular to the plane. In this case, the shadow of line AB is line BC.



Example 10.1

Below is a cube. State the projection of the following lines on plane ABCD.

- (a) AH (b) AE (c) BE (d) BH




Solution

- (a) AC (b) AD (c) BD (d) BC

Exercise 10.1

The figure below is a right pyramid with a square base, ABCD. Point O is the centre of the square.



Use the figure to answer the following questions.

1. State the projection of the following in plane ABCD.
 - (a) VA (b) VB (c) VC (d) VD
2. Name the lines that are perpendicular to VO.
3. Name the triangular planes that are perpendicular to each other.
4. State the projection of:
 - (a) line VA on plane VDB
 - (b) line VB on plane VAC.
 - (c) line VC on plane VDB.


An angle between two lines

The figure below is a cuboid.

Thus, $\tan \angle VRQ = \frac{6}{8} = 0.7500$.

Therefore, $\angle VRQ = 36.9^\circ$.

- (b) SQ and QR are sides of the right-angled triangle SQR. The angle between lines SQ and QR is SQR.




Thus, $\tan \hat{SQR} = \frac{4}{8} = 0.5000$.

Therefore, $\hat{SQR} = 26.6^\circ$

Skew lines

Lines that are not parallel and do not meet are called skew lines. The figure below is a cuboid.




From the figure, lines AH and EF are skew. They are not parallel but do not meet.

Name other skew lines from the figure.

To find the angle formed by skew lines, one of the lines must be translated. For example, from the figure above, the angle between lines FB and EH is found by translating line FB onto EA on plane HADE. When this is done, point F coincides with point E and point B with A. Thus, the angle between the two lines is \hat{HEA} .

Example 10.3

The figure below is a cuboid. The dimensions of the cuboid are 10 cm by 5 cm by 3 cm.




Find the angle between:

- (a) lines CG and DE.
- (b) Lines FG and DB.

Solutions

- (a) CG and DE are skew lines. To find the angle between them we translate DE onto CH on plane BGHC. We consider triangle GCH which is right-angled at H. The required angle is \hat{GCH}




Thus, $\tan \hat{GCH} = \frac{3}{5} = 0.6000$

Therefore, $\hat{GCH} = 31^\circ$

- (b) Lines FG and DB are skew lines. To find the angle between them, we translate DB onto plane EFGH. Point D coincides with point E and point B coincides with point G.

The angle between the lines FG and DB is formed at G, that is, \hat{FGE} . Considering the right-angled triangle EFH, EF = 3 cm and FG = 10 cm.




Thus, $\tan \hat{FGE} = \frac{3}{10} = 0.3000$.

Therefore, $\hat{FGE} = 16.7^\circ$.


Exercise 10.2

1. The figure below is a cuboid whose dimensions are 5 cm by 4 cm by 3 cm.




Find the angle between:

- lines AH and AG.
 - Lines AE and AG
 - Lines BH and AD.
2. The figure below is a cone whose diameter is 28 cm and its height is 28 cm.



Find the angle between the diameter and the line VC on the curved surface of the cone.


3. The figure below is a pyramid. The base of the pyramid is a rectangle whose dimensions are 8 cm by 6 cm. Its slanting heights are 7 cm each.



Find the angles between:

- (a) lines VQ and QS.
- (b) Lines SQ and PQ.
- (c) Lines QR and VP.

4. The figure below is a triangular prism. Its dimensions are indicated on the figure.



Find the angle between:


- (a) lines AE and AD.
- (b) Lines EB and EC.

The length of a line on a solid

To find the length of a line in three dimensions, we identify the plane containing the line. The plane may be a triangle, rectangle, parallelogram or any other plane shape. We then apply Pythagoras' theorem or trigonometric ratios to find the lengths of the lines.

Example 10.4


The figure given is a cuboid whose dimensions are 6 cm by 8 cm by 10 cm.



Find the length of line DG.

Solution


Consider $\triangle HDC$ and $\triangle GDH$ which are right angled triangles. $\triangle HDC$ is right angled at C.



$$DH^2 = 8^2 + 10^2 = 164$$

$$DH = \sqrt{164} = 12.81 \text{ cm (to 2 d.p.)}$$

Triangle GDH is right-angled at H.




$$DG^2 = DH^2 + HG^2 \\ = 164 + 36 = 200$$

$$DG = \sqrt{200} = 10\sqrt{2} \text{ cm.} \\ = 14.14 \text{ cm (to 2 d.p.)}$$


Exercise 10.3

1. The figure below is a cube of side 5 cm.




Find the length of line: (a) CE, (b) CH, (c) DH


2. The figure below is a square base pyramid. The faces of the pyramid are isosceles triangles.



- (a) Find the length of the diagonals of the base.
 (b) Find the height of the pyramid.
3. In the cuboid shown below, find the length of the line:
 (a) DG (b) GC



4. The figure below is a cuboid.



Find the length of line:


- (a) BF (b) BE.

Angle between a line and a plane

The angle between a line and a plane is the angle between the line and its projection on the plane.

Example 10.5

Below is a cuboid.




State the projection of line BE on plane BCFG and the angle between line BE and plane BCFG.

Solution

The projection of line BE on plane BCFG is BF. Thus, the angle between line BE and plane BCFG is \hat{EBF} .

Example 10.6

The figure below is a pyramid VABCD. V is the vertex of the pyramid and ABCD is the rectangular base of the pyramid. $AB = 6$ cm, $BC = 4$ cm and the slant height, $VC = 8$ cm. O is the intersection of the diagonals of the base.




- Find the angle between VA and the base.
- If M is the midpoint of AD, find the angle between VM and plane ABCD

Solution

- (a) AO is the projection of VA on the plane ABCD. In triangle VAO, the angle between line VA and plane ABCD is \widehat{VAO} .

$$\begin{aligned}AO &= \frac{1}{2} AC = \frac{1}{2} \sqrt{(36+16)} \\ &= \frac{1}{2} \sqrt{52} = \sqrt{13} = 3.61 \text{ cm.}\end{aligned}$$




$$\cos \widehat{VAO} = \frac{\sqrt{13}}{8} \approx 0.4507$$

From mathematical tables or calculator, we find an angle whose cosine is 0.4507.

$$\widehat{VAO} = 63.21^\circ \text{ (to 2 d.p.)}$$

- (b) The angle between line VN and the plane ABCD is \widehat{VMO} . The projection of line VM on plane ABCD is MO.



$$VM^2 = MO^2 + VO^2$$


$$\begin{aligned}\text{From triangle VAO above, } VO &= \sqrt{VA^2 - AO^2} \\ &= \sqrt{64 - 13} \\ &= \sqrt{51}\end{aligned}$$

$$\tan \widehat{VMO} = \frac{\sqrt{51}}{3} = 2.380$$

$$\widehat{VMO} = 67.21^\circ \text{ (to 2 d.p.)}$$


Exercise 10.4

1. The figure below is a square based pyramid. The triangular faces of the pyramid are isosceles triangles. The lengths of the slanting edges are 10 cm each.




- Find the angle between line VA and plane ABCD.
- If M is the midpoint of BC, find the angle between VM and plane ABCD.

2. The figure below is a cube of side 8 cm.




- Name the projection of line QS on plane QRUV.
- Name any two pairs of skew lines.
- Find the angle between lines RT and WV.

3. The figure below is a triangular prism.




- Calculate the length of line: (i) AC (ii) AE (iii) AD
- Calculate the angle between lines FC and FB.

4. The figure below is a cuboid.




- (a) Calculate the length of line:
- (i) BF
 - (ii) FH
 - (iii) BH
 - (iv) CH
- (b) Calculate the angles between lines:
- (i) BE and BF
 - (ii) BD and BG
 - (iii) GE and EC.

5. The figure below is a triangular prism.



Calculate:


- (a) the length of line CE
 - (b) the angle between lines CE and EA.
6. The figure below is a cuboid.



- (a) Name the projection of line BE on plane ABGH


- (b) Find the length of line:
 (i) BF (ii) BE
 (c) Find the angle between lines BE and BF.

7. The figure below is a cube.



- (a) Name:
 (i) the projection of line QS on plane QRUV.
 (ii) The projection of line RV on plane RSTU.
 (b) Find the angle between lines QT and WQ.
 (c) Calculate the angle between line QS and plane QRUV.

8. The figure below is a square based pyramid with vertex V. Point O is the intersection of diagonals AC and BD.



- (a) Name the projection of line VB on plane ABCD.
 (b) Find the angle between line VA and line AC
 (c) Calculate the angle between line VB and plane ABCD.


9. The figure below is a triangular prism. Its base is a right-angled triangle.

- (a) Calculate the length of line:
 (i) QS (ii) UQ (iii) UR
 (b) Calculate:
 (i) \hat{URS} (ii) \hat{UQS} .

Angle between two planes

The angle between two intersecting planes is the angle between two lines, one in each plane, such that the two lines meet at a point on the line common to the planes, and both lines are at right angles to the common line.


The diagram below shows two intersecting planes ABCD and BCEF.



The line of intersection of the two planes is BC. Lines MN and NP are perpendicular to line BC at N. The angle between the two planes is \hat{MNP} .


Example 10.7

The figure below is a cuboid. Find the angle between planes BCFG and ACFH.



Solution

The angle between the two planes is \hat{ACB} .




From triangle ABC,

$$\tan \hat{ACB} = \frac{5}{7} \approx 0.7143$$

$$\hat{A}CB = 35.54^\circ$$


Exercise 10.5

1. The figure below is a cuboid.




From the figure find the angles between planes:

- ABGF and AFHC.
 - BGED and DEHC.
 - ABCD and BHEA.
 - DEGB and BGHC.
2. The figure below shows a right pyramid on a square base. The slanting edges are 8 cm long each.




Find:

- the height of the pyramid.
 - the angle the slanting edges make with the base.
3. The following figure is a triangular prism.
- Find:
- height CD.
 - the angle between lines AC and AD.




4. The following figure shows a square based pyramid of side 6 cm. Point N is the midpoint of BC. The slanting edges are 5 cm each.



Find:


- (a) the height of the pyramid.
 - (b) The angle between planes VBC and ABCD.
5. The faces of a square based pyramid are isosceles triangles. The sides of the square base are 12 cm. The lengths of the slanting edges are 10 cm each.
- (a) Calculate the height of the pyramid.
 - (b) Find the angle between the slanting edges and the base.
6. The height of a circular cone is 35 cm and the radius of the base is 21 cm.
- Find:
- (a) the length of the slanting height.
 - (b) the angle between the slanting height and the base.
7. A pyramid has a rectangular base, 12 cm long and 9 cm wide. Its slanting faces are isosceles triangles whose edges are 15 cm long. Find:
- (a) the height of the pyramid.
 - (b) the angle between a slanting edge and the base.
 - (c) the angle between the triangular faces and the base.

8. The figure below is a pyramid on a rectangular base. $AB = 16$ cm, $BC = 12$ cm and $VA = 13$ cm.




Find:

- the length of the line BD .
 - the height of the pyramid
 - the angle between VB and the base.
 - the angle between plane VBC and the base.
9. The figure below is a tetrahedron. The base is an equilateral triangle of side 6 cm. Point N is the midpoint of CD . The slanting edges of the tetrahedron are 10 cm each. AO is the height of the tetrahedron.




Find:

- the height of the tetrahedron.
 - the angle between line AD and plane BCD .
 - the angle between planes ACD and BCD .
10. The following figure is a tetrahedron in which $AB = BC = AC = 8$ cm and $AD = DC = BD = 10$ cm. Find the angle between planes ABC and DBC .




11. The following figure is a right pyramid with a rectangular base, PQRS, and vertex V. $PQ = 4$ cm and $QR = 6$ cm. The slanting edges of the pyramid are 8 cm each. Point M is the midpoint of QR.



Calculate:


- the length of line VO.
 - the angle between lines VS and SQ.
 - the angle between line VM and plane PQRS.
12. The figure below is a triangular prism. The cross section of the prism is an isosceles triangle.



Calculate:


- the angle between lines AC and BC.
- The angle between lines CF and CD.
- The angle between planes ABFE and ACDE.

13. The figure below is a cube. Points M and N are the midpoints of YV and XW respectively.




Find:

- the length of line SN.
 - the angle between planes MNSR and UTSR.
 - the angle between planes VWSR and MNSR.
 - the angle between VU and NS.
14. The figure below is a wedge. $TU = 16$ cm, $QR = 8$ cm and $RU = 5$ cm.



- Write down the projection of line TQ onto plane PQRS.
 - Find the angle between line TQ and plane PQRS.
 - Find the angle between planes PQUT and STUR.
15. The figure below represents a roof of a house.




Find:

- the height of the roof,


(b) the angle between planes AMJ and MJKL.

16. The figure below is a right pyramid, VABCD. The base is a square, 10 cm each side. The vertex is vertically above the centre of the square. $VA = VB = VC = VD = 9$ cm. Points P and N are the midpoints of VO and AB respectively.




Find:

- the length of the line VO.
 - the angle between lines VA and AC.
 - the projection of line PN on plane ABCD
 - the angle between line PN and plane ABCD.
 - the angle between planes VBC and ABCD.
17. The figure below is a pyramid with a square base, ABCD, of length 5 cm. Edge VA is 5 cm long and perpendicular to the base at A.



- Name the projection of line VC on plane ABCD.
 - Find the length of edge VC.
 - Find the angle between edge VC and plane ABCD.
18. The following figure is a cone whose base radius is 14 cm. The perpendicular height, VO, is 18 cm. Point P is such that


$$OP = \frac{2}{3}OV.$$




- (a) Find the lengths of lines VB and PB.
- (b) Find the angle between lines:
 - (i) VB and BA.
 - (ii) AP and AO.

19. The following diagram shows a frustum of a square based pyramid. The base, ABCD, is a square of side 10 cm long. The top, PQRS, is a square, 4 cm a side, and each of the short edges of the frustum is 5 cm. Determine:

- (a) the height of the frustum.
- (b) The angle between AR and the base, ABCD.



20.



The figure represents a solid frustum cut off from a cone of height 4.2 cm. The radii of the circular ends are 1.4 cm and 0.7 cm. The slanting height of the frustum is 10 cm.

Calculate:

- (a) the volume of the frustum.
- (b) the surface area of the frustum.

11. Further Logarithms

Logarithmic notation

When a number is expressed in the index form a^x , x is called the logarithm and a is called the base.

The logarithm of y to the base a is written as $\log_a y$, i.e. if $y = a^x$, then

$$\log_a y = x.$$

$$1000 = 10^3, \text{ therefore, } \log_{10} 1000 = 3.$$

$$625 = 5^4, \text{ therefore, } \log_5 625 = 4.$$

Similarly, if $p = \log_r q$, then, $q = r^p$.

Laws of logarithms

The three laws of logarithms are:

(i) $\log_x(AB) = \log_x A + \log_x B.$

(ii) $\log_x(A \div B) = \log_x A - \log_x B.$

(iii) $\log_x A^n = n \log_x A.$

The above three laws are derived from the rules of indices.

$$x^m \times x^n = x^{(m+n)}, x^m \div x^n = x^{m-n} \text{ and } (x^m)^n = x^{mn}$$

Let $x^m = A$ and $x^n = B$, then

$$\log_x A = m, \text{ and } \log_x B = n.$$

(i) Now $AB = x^m \times x^n$

$$\text{Therefore, } AB = x^{m+n}$$

$$\text{So, } \log_x(AB) = m + n$$

$$\therefore \log_x(AB) = \log_x A + \log_x B$$

(ii) $\frac{A}{B} = \frac{x^m}{x^n}$. Therefore, $\frac{A}{B} = x^{m-n}$

$$\text{So, } \log_x \frac{A}{B} = m - n$$

$$\text{Therefore, } \log_x \frac{A}{B} = \log_x A - \log_x B.$$

(iii) $(x^m)^n = x^{mn}$

$$\text{Let } A = x^m \text{ such that } \log_x A = m, \text{ then } A^n = x^{nm} = x^{mn}$$

$$\log_x(A^n) = mn = nm = n \log_x A \text{ since } m = \log_x A.$$

In each of these laws every logarithm must be to the same base.

Example 11.1

$$(a) \log_x(35.7 \times 40.3) = \log_x 35.7 + \log_x 40.3$$

$$(b) \log\left(\frac{8.1}{9.37}\right) = \log 8.1 - \log 9.37$$

$$(c) \log_x(39.3^4) = 4 \log_x 39.3$$

$$(d) \log_x \sqrt{65.7} = \log_x 65.7^{\frac{1}{2}} = \frac{1}{2} \log_x 65.7.$$

In general,

$$\log_x \sqrt{a} = \frac{1}{2} \log_x a, \log_x \sqrt[3]{a} = \frac{1}{3} \log_x a, \log_x \sqrt[n]{a} = \frac{1}{n} \log_x a$$

Logarithms to the base 10, are abbreviated as $\log x$. When the base is not given, it is generally taken to be base 10.

Logarithm of unity

For any non-zero number a , $a^0 = 1$,
Therefore, $\log_a 1 = 0$.

Logarithm of the base

Since $a^1 = a$, then $\log_a a = 1$.

For example, $\log_{10} 10 = 1$, $\log_5 5 = 1$, $\log_{32} 32 = 1$.

$\log_a a^5 = 5 \log_a a = 5(1) = 5$

In general, $\log_a a^n = n \log_a a = n$.

Simplification of logarithmic expressions

Example 11.2

Simplify the following without using tables:

$$(a) \begin{aligned} \log 8 + \log 12.5, \\ = \log(8 \times 12.5) \\ = \log 100 \\ = 2 \end{aligned}$$

$$(b) \begin{aligned} \log 40 - \log 4, \\ = \log(40 \div 4) \\ = \log 10 \\ = 1 \end{aligned}$$

$$(c) \begin{aligned} \frac{\log 27}{\log 3} &= \frac{\log 3^3}{\log 3} \\ &= \frac{3 \log 3}{\log 3} \\ &= 3. \end{aligned}$$

Example 11.3



If $\log 2 \approx 0.30103$ and $\log 3 \approx 0.47712$, find without using tables of calculators, correct to 4 decimal places:

$$\begin{aligned} \text{(a) } \log 6 &= \log (2 \times 3) \\ &= \log 2 + \log 3 \\ &= 0.30103 + 0.47712 \\ &= 0.77815 \approx \mathbf{0.7782} \end{aligned}$$

$$\begin{aligned} \text{(b) } \log (30) &= \log (3 \times 10) \\ &= \log 3 + \log 10 \\ &= 0.47712 + 1 \\ &= 1.47712 \approx \mathbf{1.4771} \end{aligned}$$

$$\begin{aligned} \text{(c) } \log 72 &= \log (8 \times 9) \\ &= \log 8 + \log 9 \\ &= \log 2^3 + \log 3^2 \\ &= 3\log 2 + 2\log 3 \\ &= 3 \times 0.30103 + 2 \times 0.47712 \\ &= 0.90309 + 0.95424 \\ &= 1.85733 \approx \mathbf{1.8573}. \end{aligned}$$

$$\begin{aligned} \text{(d) } \log 5 &= \log\left(\frac{10}{2}\right) \\ &= \log 10 - \log 2 \\ &= 1 - 0.30103 \\ &= \mathbf{0.69897}. \end{aligned}$$

Example 11.4

If $\log x = p$ and $\log y = q$, express in terms of p and q :

$$\text{(a) } \log (x^2y)$$

$$\text{(b) } \log\left(\frac{x^3}{y^4}\right)$$

$$\text{(c) } \log \frac{\sqrt{x^3}}{y},$$

$$\text{(d) } \log \frac{x^2}{10y}.$$

Solution

$$\begin{aligned} \text{(a) } \log(x^2y) &= \log x^2 + \log y \\ &= 2 \times \log x + \log y \\ &= \mathbf{2p + q}. \end{aligned}$$

$$\begin{aligned} \text{(b) } \log\left(\frac{x^3}{y^4}\right) &= \log x^3 - \log y^4 \\ &= 3 \times \log x - 4 \times \log y \\ &= \mathbf{3p - 4q}. \end{aligned}$$

$$\begin{aligned} \text{(c) } \log \frac{\sqrt{x^3}}{y} &= \log \sqrt{x^3} - \log y \\ &= \log x^{\frac{3}{2}} - \log y \\ &= \frac{3}{2} \log x - \log y \end{aligned}$$

$$\begin{aligned} \text{(d) } \log \frac{x^2}{10y} &= \log x^2 - (\log 10y) \\ &= 2 \log x - (\log 10 + \log y) \\ &= \mathbf{2p - (1 + q)} \end{aligned}$$

$$= \frac{3}{2}p - q.$$

$$= 2p - q - 1.$$

Exercise 11.1

In all questions consider base 10

Express in the form $\log x$ and then simplify where possible (Q. 1 - Q. 6):

- | | |
|---------------------------------|-----------------------------------|
| 1. $\log 2 + \log 5.$ | 2. $2\log 5 + \frac{1}{2}\log 16$ |
| 3. $\log 12 + \log 25 - \log 3$ | 4. $2\log p + \log q$ |
| 5. $\log 12.5 + 3\log 2$ | 6. $3\log x + \frac{1}{2}\log y$ |

Write each of the following as the sum or difference of two logarithms (Q. 7 - 9):

- | | |
|--------------------------|--------------------------------|
| 7. $\log(a^3b^{-2})$ | 8. $\log \frac{\sqrt{x}}{y^2}$ |
| 9. $\log(\sqrt[3]{x^2})$ | |

Simplify, without using tables or calculators, Q. 10 - 17:

- | | |
|---|---|
| 10. $\frac{\log 4}{\log 8}$ | 11. $\frac{\log 27}{\log 81}$ |
| 12. $\frac{\log 1000}{\log 100}$ | 13. $\log 50 - \log 100 + \log 2$ |
| 14. $\log 54 - \log 15 + 2 \log \left(\frac{5}{3}\right)$ | 15. $3 \log 5 + \log 10 - \log 625 - \log 2.$ |
| 16. $\log 120 + \frac{1}{3}\log 27 - 2 \log 6$ | 17. $\frac{\log 32}{\log 49} \frac{\log 8}{\log 7} \frac{\log 0.8}{\log 0.7}$ |
18. If $\log_a x = 2.526$, find the values of:
- | | |
|-----------------------|------------------------------------|
| (a) $\log_a(x^{10}),$ | (b) $\log_a(\sqrt{x}),$ |
| (c) $\log_a(a^3x^2),$ | (d) $\log_a \frac{a^2}{\sqrt{x}}.$ |
19. If $\log_{10} 2 = 0.301030$ and $\log_{10} 3 = 0.477121$, find without using calculators or tables, correct to 5 d.p. the values of:

(a) $\log_{10} 18$,

(b) $\log_{10} 1.5$,

(c) $\log_{10} 8$,

(d) $\log_{10} 4.5$,

(e) $\log_{10} \sqrt[3]{60}$.

20. If $\log x = a$, $\log y = b$ and $\log z = c$, express the following in terms of a , b and c .

(a) $\log(x^2)$,

(b) $\log(y^3)$,

(c) $\log(\sqrt{z})$,

(d) $\log \frac{x^2 y}{\sqrt{z}}$.

(e) $\log(xy^2)$,

(f) $\log \frac{x}{y^3}$,

(g) $\log \sqrt{\frac{10x^2}{y}}$,

(h) $\log \sqrt[3]{\frac{100x^3}{y^2}}$.

Solution of logarithmic equations

The equations involving $\log x$ terms may be solved using the laws of logarithms.

Example 11.5

Solve the equation: $\log(7x + 2) - \log(x - 1) = 1$.

Solution

$\log(7x + 2) - \log(x - 1) = \log 10$, since $\log 10 = 1$

$\Leftrightarrow \log \frac{7x + 2}{x - 1} = \log 10$. Hence, $\frac{7x + 2}{x - 1} = 10$

$$7x + 2 = 10x - 10$$

$$12 = 3x$$

$$x = 4.$$

The equation $a^x = b$

Example 11.6

Solve the equations:

(a) $3^{2x-1} = 2187$

(b) $2^x = 5$

Solution

(a) $3^{2x-1} = 2187$

$3^{2x-1} = 3^7$

$\therefore 2x - 1 = 7$

$2x = 8$

$x = 4$

(b) $2^x = 5$ Taking logarithm on both sides, $\log(2^x) = \log 5$

$$\text{Therefore, } x \log 2 = \log 5 \Rightarrow x = \frac{\log 5}{\log 2} = \frac{0.6990}{0.3010} \approx 2.322$$

Example 11.7

Solve the equation: $5^{2x+2} = 26^{x-1}$.

Solution

$$5^{2x+2} = 26^{x-1}$$

$$(5^2)^x \times 5^2 = (26)^x \times 2^{-1}$$

$$25^x \times 25 = 64^x \times \frac{1}{2} \text{ multiplying by 2 on both sides gives,}$$

$$50(25^x) = 64^x$$

$$50 = \left(\frac{64}{25}\right)^x = 2.56^x$$

Taking logs on both sides

$$\log(50) = \log(2.56^x) = x \log 2.56$$

$$\text{Therefore, } x = \frac{\log 50}{\log 2.56} \approx 4.162$$

Alternatively:

$$\log(5^{2x+2}) = \log(26^{x-1})$$

$$(2x+2) \log 5 = (x-1) \log 26$$

$$2x \log 5 + 2 \log 5 = x \log 26 - \log 26$$

$$2 \log 5 + \log 26 = x(\log 26 - 2 \log 5)$$

$$\log 5^2 + \log 26 = x(\log 26 - 2 \log 5)$$

$$\log(5^2 \times 26) = x(\log 26^2 - \log 5^2)$$

$$\log 50 = x \log\left(\frac{64}{25}\right) = x \log 2.56$$

$$\text{Therefore, } x = \frac{\log 50}{\log 2.56} \approx 4.162$$

Example 11.8

Solve the equation: $2^{2x+3} - 7(2^{x+1}) - 15 = 0$

Solution

$$2^{2x+3} = 2^{2x} \times 2^3 = (2^x)^2 \times 8 = 8(2^x)^2$$

$$2^{x+1} = 2^x \times 2^1 = 2^x \times 2 = 2(2^x)$$

$$\text{Therefore, } 2^{2x+3} - 7(2^{x+1}) - 15 = 0 \Rightarrow 8(2^x)^2 - 7 \times 2(2^x) - 15 = 0$$

Let $2^x = y$,

$$8y^2 - 14y - 15 = 0$$

Factorizing, we get $(2y - 5)(4y + 3) = 0$

$$y = \frac{5}{2} \text{ or } y = -\frac{3}{4}$$

Therefore, $2^x = 2.5$ or $2^x = -0.75$ (not possible as 2^x is always positive)

Taking logs on both sides,

$$\log 2^x = \log 2.5$$

$$x \log 2 = \log 2.5$$

$$x = \frac{\log 2.5}{\log 2} \approx 1.322.$$

Exercise 11.2

Solve the equations: (Q. 1 - 13)

1. $\log (3x + 4) = \log (x - 2) + 3 \log 2$
2. $\log (3y - 1) = \log (2y + 1) - \log 4$
3. $\log_{10} (3x - 5) = 2 \log_{10} 5 - 1$
4. $\log_{10} (10y + 5) - \log_{10} (y - 4) = 2$
5. $\log_{10} (9x + 1) = 2$
6. $\log_{10} (8x^3) = 3$
7. $\log (x - 1) + \log x = \log 20$
8. $2 \log x = \log (x + 6)$
9. $2 \log x - \log (2x + 3) = 0$
10. $\log (x^2 + 2) - \log (3x + 6) = 0$
11. $2 \log (x - 3) - \log (x + 17) = 0$
12. $1 + 2 \log_{10} x - \log_{10} (x^2 - 7x + 2) = 0$
13. $\log (x^2 - 8x - 17) = \log 4 + \log (x - 1)$
14. If $\log a^4 - \log a^2b + \log b^2 = 0$, express b in terms of a .
15. Find x , if $\log_x 2 + \log_x 8 = 4$
16. Find y , if $\log 5 + \log y = 0$.
17. Find z , if $4 \log_z 3 + 2 \log_z 2 - \log_z 144 = 2$
18. Solve the equations, giving your answers to 4 S.F:

(a) $3^x = 5,$	(b) $2^{x+1} = 3,$
(c) $2^x = 3^{x+2},$	(d) $5^{2x+1} = 3^{2-x},$

19. Expressing the numbers as powers of 2 and using the fact that $\log_a a = 1$, find the values of:

- (a) $\log_2 8$,
(b) $\log_2 \frac{1}{4}$,
(c) $\log_2 64$.

20. Solve the equations:

- (a) $\log_2 x + \log_2 (x - 30) = 6$,
(b) $2(\log_{10} x - \log_{10} 4) = 1 - \log_{10} 5$,
(c) $\log(2x - 11) - \log 3 = \log 2 - \log x$.

21. Find x, if $2\log 4 - \log x = 2\log \frac{4}{7} + \log \frac{7}{2} - \log \frac{5}{14}$

22. Solve the equations:

- (a) $4^{x+1} = 128$,
(b) $8^x = 32$,
(c) $3^{2x} \cdot 3^{x-1} = 81$

23. Solve the equations:

- (a) $3^x \cdot 2^{2x-3} = 18$,
(b) $4^{2x+1} = 2^{x+4}$,
(c) $3^{2x+1} - 14(3^x) - 5 = 0$
(d) $3(2^x) = 2^{4-2x}$.

24. Solve the following equation, expressing in terms of 2^x , 3^x , ...

- (a) $2^{2x+1} - 9(2^x) + 4 = 0$,
(b) $3(3^{2x}) - 28(3^x) + 9 = 0$
(c) $4^x - 10(2^x) + 16 = 0$.

25. Using the substitution $y = 3^x$, solve the equation $4(3^{2x+1}) - 17(3^x) - 7 = 0$.

26. Without using tables, evaluate:

$$\log \frac{41}{35} + \log 70 - \log \frac{41}{2} + 2 \log 5.$$